


**Report on Propionic Acid, Sodium Propionate, Calcium Propionate, Thiodipropionic  
Acid & Dilauryl Thiodipropionic Acid**

**1/28/74**

**Q 30**



FOOD AND DRUG  
LABORATORIES, INC.

VOL. I

Q 30

R E P O R T

PROPTONIC ACID

CAS. REG. NO. 000079-09-4

SODIUM PROPIONATE

CAS. REG. NO. 000137-40-6

CALCIUM PROPIONATE

CAS. REG. NO. 004075-81-4

THIODIPROPIONIC ACID

CAS. REG. NO. 000111-17-1

DILAURYL THIODIPROPIONIC ACID

CAS. REG. NO. 000123-28-4

Submitted to:


GRAS Review Branch (BF-335)  
Bureau of Foods  
Food and Drug Administration  
200 C Street, S. W.  
Washington D.C., 20204


Attention:

Mr. Alan Spiher,  
Project Manager

Date: January 28, 1974

Laboratory No: 1216

  
James Bond, Manager  
Monograph Group

  
Howard Feiman, Director  
Biological Sciences

## TABLE OF CONTENTS

<u>SUBJECT</u>	<u>PAGE</u>
Summary	1
Chemical Information	5
Acute Toxicity	21
Short-term Studies	23
Long-term Studies	26
Special Studies	28
<u>Bibliography</u>	
Propionic Acid	B-1
Sodium Propionate	B-34
Calcium Propionate	B-41
Thiodipropionic Acid	B-48
Dilauryl Thiodipropionic Acid	B-50

## SUMMARY

In this review propionic acid, sodium propionate, calcium propionate, thiodipropionic acid, and dilauryl thiodipropionate are considered. Propionic acid is a member of the aliphatic monocarboxylic acid series; it is a liquid, has a strong odor and is somewhat corrosive. The propionates used in the food industry as antimicrobials are the sodium and calcium salts, which yield free acid in the pH range of the foods in which they are used.

Under federal regulation, propionates are generally recognized as safe for use in foods and no upper limits are imposed on usage, excepting bread, rolls, and cheese which come under standard of identity. Sodium and calcium propionate are limited to 0.32% of flour in white bread and rolls and to 0.38% in the corresponding whole wheat products; in cheese products they are limited to 0.3% of the cheese. The propionates are not mentioned in the food additive regulations of France, West Germany, Greece, Portugal, and Turkey but they are included in those of other western European countries for which information concerning them is at hand. The major usages of propionates are to prevent mold and rope in baked goods and mold in certain types of cheese (407a).

Propionic acid is used as an esterifying agent in the production of cellulose, other propionates are used as mold inhibitors, in the manufacture of ester solvents, fruit flavors, and perfume bases (234a).

Sodium propionate has been used as a topical antifungal agent and in chronic bacterial conjunctivitis and blepharitis. In the veterinary field, it is used in the prevention of bovine ketosis, dermatosis, wound infections, conjunctivitis, vaginitis, and otitis (234a).

Calcium propionate has been used as an inhibitor of molds and other micro-organisms in foods, tobacco, and in the production of butyl rubber to improve processibility and scorch resistance, while medically it has been used as an antifungal agent in dermatology (234a).

Thiodipropionic acid is used as an antioxidant for soap products, as an esterifier in plasticizers and lubricants and is proposed for use in edible fats, oils and other foods (234a).

The acute toxicities for various species and for various routes of administration are shown in Table 1 of the text for propionic acid, sodium propionate, calcium propionate, thiodipropionic acid and di-lauryl thiodipropionate. In acute rabbit eye irritation studies, the edema produced by propionate was quite similar to the edema produced with the same molar concentration of hydrochloric acid (196).

The addition of propionate to diets fed to varying species at levels of 24% or below did not affect growth rate in the adult animals but did affect the growth rate and physical well-being of weanling rats fed a diet containing 24% (231, 461, 571, 85).

When sodium propionate was fed to rats as 75% of the diet for 32 weeks, a depression of growth during the first week of feeding was observed, but it did not influence mortality and did not produce any detectable effects on hemoglobin levels, organ weights or histopathology of the tissues (412).

A diet containing 10% sodium propionate fed twice daily for 50 days increased the amount of fat in sheep at slaughter, indicating that they converted digestible energy to weight gains considerably better than did the controls (85).

In vitro studies utilizing human skin and fetal lung cells and strain L cells showed that a concentration of 10 mg% sodium propionate was toxic to these cell lines, but 1 mg% had no significant effect. Propionic acid and thiodipropionic acid showed a great anticarcinogenic activity against the growth of transplanted tumor cells in mice (347, 571).

In intestinal sections, propionic acid has an inhibitory effect on the transfer of fatty acids; the concentrations necessary to produce the inhibition decrease with the increase in length of the fatty acid chain. In the ingesta of chickens, fatty acids are completely removed from the ingesta before it reaches Meckel's diverticulum; in mice the preferential movement of propionic acid is in the jejunum rather than in the ileum (55, 35).

In Vitamin B<sub>12</sub>-deficient humans, the injected radioactivity of propionates found in urinary methylmalonic acid is small; 20 to 50% of the propionate administered is the daily turnover in normal patients and 20% in Vitamin B<sub>12</sub>-deficient patients. The fate of the remainder is unknown. Studies show that propionic acid is the second most effective of the fatty acids in penetrating the red blood cells, thus producing hemolysis, and that increasing the osmotic concentration of the outside fluid retards hemolysis.

The electroencephalogram patterns of the adult rabbits which were injected i.v. with a propionate solution showed changes in the EEG pattern from a resting type to one of sleep. These patterns were very similar to those observed under deep sodium penabarbato1 anesthesia (372).

## CHEMICAL INFORMATION

### I. Nomenclature

#### A. Common names (234a)

##### Propionic acid-

##### Sodium propionate-

propionic acid sodium salt

##### Calcium propionate-

propionic acid calcium salt

##### Thiodipropionic acid-

No other common names were encountered in the literature searched.

##### Dilauryl thiodipropionate acid

No other common names were encountered in the literature searched.

#### B. Chemical names (234a, 176a)

##### Propionic acid-

propionic acid, methylacetic acid, ethylformic acid

##### Sodium propionate-

No other chemical name encountered in the literature searched.

##### Calcium propionate-

No other chemical name encountered in the literature searched.

##### Thiodipropionic acid-

3, 3'-thiodipropionic acid,  $\beta, \beta$ -thiodipropionic acid, thiodihydracrylic acid, diethylsulfide 2, 2'-dicarboxylic acid.

##### Dilauryl thiodipropionate acid

No other chemical name encountered in the literature searched.



C. Trade Names ( 234a, 258a)

Propionic acid-

No other trade names were encountered in the literature searched.

Sodium propionate-

Mycoban

Calcium propionate-

Mycoban

Thiodipropionic acid-

$\beta$ ,  $\beta'$ thiodipropionic acid

Dilauryl thiodipropionate acid

b, b'dilauryl thiodipropionate,  
Milban F.

D. Chemical Abstracts Services Unique Registry Number

Propionic acid

000 070-09-4

Sodium propionate

000137-40-6

Calcium propionate

004075-81-4

Thiodipropionic acid


000111-17-1

Dilauryl thiodipropionate


000123-28-4

II. Empirical Formula (234a, 105a)


Propionic acid




Sodium propionate




Calcium propionate



Thiodipropionic acid




Dilauryl thiodipropionate




III. Structural Formula (234a, 407a, 105a)


Propionic acid




Sodium propionate




Calcium propionate



Thiodipropionic acid



Dilauryl thiodipropionate



IV. Molecular Weight (234a, 105a)

Propionic acid

74.08

Sodium propionate

96.07

Calcium propionate

186.22

Thiodipropionic acid

178.20

Dilauryl thiodipropionate

514.85

V. Specifications

A. Chemical (234a)

Propionic acid

C 48.64%

H 8.16%

O 43.20%

Sodium propionate

C 37.50%

H 5.25%

O 33.31%

Na 23.94%

Propionic acid 77.11%

Calcium propionate

C	38.70%
H	5.41%
O	34.37%
Ca	21.52%

Thiodipropionic acid

C	40.44%
H	5.66%
O	35.91%
S	17.99%

Dilauryl thiodipropionate

C	70.0%
H	11.3%
O	12.4%
S	6.2%

B. Food Grade (105a)

Propionic acid

**Assay.** Not less than 99.5 percent of  $C_3H_6O_2$ .

**Distillation range.** Between  $138.5^\circ$  and  $142.5^\circ$ .

**Specific gravity.** Between 0.993 and 0.997 at  $20^\circ/20^\circ$ .

**Limits of Impurities**

**Aldehydes** (as propionaldehyde). Passes test (limit about 0.05 percent).

**Arsenic** (as As). Not more than 3 parts per million (0.0003 percent).

**Heavy metals** (as Pb). Not more than 10 parts per million (0.001 percent).

**Nonvolatile residue.** Not more than 0.01 percent.

**Readily oxidizable substances** (as formic acid). Passes test (limit about 0.05 percent).

**Water.** Not more than 0.15 percent.

### Sodium propionate

Assay. Not less than 99.0 percent of  $C_3H_5NaO_2$  after drying.

#### Limits of Impurities

Alkalinity (as  $Na_2CO_3$ ). Passes test (about 0.15 percent).

Arsenic (as As). Not more than 3 parts per million (0.0003 percent).

Heavy metals (as Pb). Not more than 10 parts per million (0.001 percent).

Iron. Not more than 30 parts per million (0.003 percent).

Water. Not more than 1 percent.

### Calcium propionate

Assay. Not less than 98.0 percent of  $C_6H_{10}CaO_4$ , calculated on the anhydrous basis.

#### Limits of Impurities

Arsenic (as As). Not more than 3 parts per million (0.0003 percent).

Fluoride. Not more than 30 parts per million (0.003 percent).

Heavy metals (as Pb). Not more than 10 parts per million (0.001 percent).

Insoluble substances. Not more than 0.2 percent.

Magnesium (as MgO). Passes test (about 0.4 percent).

Water. Not more than 5 percent.

### Thiodipropionic acid

No Food Grade information was encountered in the available literature.

### Dilauryl thiodipropionate acid

Assay. Not less than 99.0 percent of  $C_{30}H_{58}O_4S$ .

Solidification point. Not below 40°.

#### Limits of Impurities

Acidity (as thiodipropionic acid). Not more than 0.2 percent of  $C_6H_{10}O_4S$ .

Arsenic (as As). Not more than 3 parts per million (0.0003 percent).

Heavy metals (as Pb). Not more than 20 parts per million (0.002 percent).

Lead. Not more than 10 parts per million (0.001 percent).

## C. Official Compendia

Food Chemical Codex

## VI. Description

### A. General Characteristics (105a, 234a)

#### Propionic acid

An oily liquid having a slightly pungent, rancid odor. It is miscible with water and with alcohol and various other organic solvents.

#### Sodium propionate

Colorless, transparent crystals or a granular crystalline powder. It is odorless, or has a faint acetic-butyric odor. It is deliquescent in moist air.

#### Calcium propionate

White crystals or crystalline solid, possessing not more than a faint odor of propionic acid.

#### Thiodipropionic acid

Nacreous leaflets from hot water.

#### Dilauryl thiodipropionate acid

White crystalline flakes having a characteristic sweetish, ester-like odor.

### B. Physical Properties (176a, 105a, 234a)

#### Propionic acid

melting point, °C	-22
boiling point, °C	141.1
vapor pressure, mm Hg, at 28.0°C	5
at 52.0°C	20
at 85.8°C	100
at 122.0°C	400
specific gravity, 20/4°C	0.992
20/20°C	0.9952
refractive index, $n_D^{20}$	1.3874
viscosity at 25°C, cP	1.035
surface tension in air at 20°C, dyn/cm	26.70
coefficient of cubical expansion at 0-133°C/°C	$1.102 \times 10^{-3}$
critical temperature, °C	337.6
critical pressure, atm	53.0
specific heat at 20-137°C, cal/g	0.726
heat of fusion at melting point, cal/g	23.4
heat of vaporization at 139.3°C, cal/g	98.9
heat of combustion of liquid, cal/g	4931.5
flash point, Cleveland open cup, °F	150
weight at 20°C, lb/gal	8.28

Sodium propionate

One gram is soluble in about 1 ml of water at 25°, in about 0.65 ml of boiling water, and in about 24 ml of alcohol. The pH of a 1 in 10 solution is between 8.0 and 10.5.

Calcium propionate

One gram dissolves in about 3 ml of water. The pH of a 1 in 10 solution is between 8 and 10. Slightly soluble in methanol, ethanol; practically insoluble in acetone and benzene.

Thiodipropionic acid

m. p. 134°C

Kat 25°C  $7.8 \times 10^{-5}$

One gram dissolves in 26.9 ml H<sub>2</sub>O at 26°C. Freely soluble in hot water, alcohol, acetone.

Dilauryl thiopropionate

Insoluble in water.

Soluble in most organic solvents.

C. Stability in Containers \* (176a, 407a, 105a)

Propionic acid

Handle as any organic chemical of comparable flash point and vapor pressure. Uncoated steel containers are not recommended.

Sodium propionate

Emits acrid fumes at high temperatures.

Presents little hazard in storage.

Calcium propionate

Emits acrid fumes at high temperature, protect from humidity, indefinite shelf-life.

Thiodipropionic acid

No stability information encountered in the literature searched.

Dilauryl thiopropionate

Store in well-closed containers.

---

\* Private communication from unidentified source provided by the FDA.

## VII. Analytical

Compound:	Propionic acid
Title:	Gas chromatographic determination of propionic, sorbic, and benzoic acids in rye bread and margarine (123a).
Principle determination step:	Flame ionization detector
Principle separation technique:	Extraction and mixed carbowax, terephthalic acid on chromosorb W(AW-DMCS).
Principle analytical reagent:	Valeric acid
Moiety measured:	Unknown



Compound:	Sodium propionate
Title:	Separation and identification of sodium salts of acetic, propionic, butyric and valeric acids by paper chromatography (475).
Principle determination step:	Color
Principle separation technique:	Silicic acid column
Principle analytical reagent:	Methyl red, bromothymol blue in formalin at pH 5.2.
Moiety measured:	Acid group.

Compound:	Propionic acid
Title:	Determination of volatile free fatty acids of human blood (221).
Principle determination step:	Flame ionization
Principle separation technique:	Extraction, gas liquid chromatography.
Principle analytical reagent:	Chloroform-methanol-sulfuric acid mixture.
Moiety measured:	Unknown

Compound:	Propionic acid
Title:	Identification of acetic, propionic and sorbic acid in bakery products by thin layer chromatography (346).
Principle determination step:	Color
Principle separation technique:	Distillation, thin layer chromatography.
Principle analytical reagent:	H <sub>3</sub> PO <sub>4</sub> , bromcresol purple.
Moiety measured:	Acid group

Compound:	Thiodipropionic acid
Title:	The multi-determination of antioxidants in lard (567).
Principle determination step:	Infrared spectrophotometry
Principle separation technique:	Vacuum sublimation, gas chromatography.
Principle analytical reagent:	Diazomethane
Moiety measured:	All IR absorbing bonds.

Compound:	Dilauryl thiodipropionate
Title:	Determination of DLTDPP and other antioxidants by a modified sublimation technique (583).
Principle determination step:	Spectrophotometry
Principle separation technique:	Sublimation, gas liquid chromatography
Principle analytical reagent:	1% Apiezon N on gas Chrom Q.
Moiety measured:	All IR absorbing bonds.

Calcium propionate: No method specifically for calcium propionate was found in the literature searched. However, the procedure of Young et al (475) implies use for propionates in general.

## VIII. Occurrence and Levels Found in:

### A. Plants (248a, 234a)

#### Propionic acid

Apples, strawberries, tea, violet leaves.  
Wood distillate. Bacterial action on many materials.

#### Sodium propionate

No occurrences or levels found in plants encountered in the literature searched.

#### Calcium propionate

No occurrences or levels found in plants encountered in the literature searched.

#### Thiodipropionic acid

No occurrences or levels found in plants encountered in the literature searched.

#### Dilauryl thiodipropionate

No occurrences or levels found in plants encountered in the literature searched.

### B. Animals (176a, 234a, 407a)

#### Propionic acid

Stomach of ruminants  
Dairy products - small amounts

#### Sodium propionate

No occurrences or levels found in animals encountered in the literature searched.

#### Calcium propionate

No occurrences or levels found in animals encountered in the literature searched.

# I Acute Toxicity

Table 1

Substance	Animal and Strain	Number	Route	LD <sub>50</sub> (mg/kg body wt.)	Ref.
Propionic acid	Mice	Minimum 6 groups x 10	i. v.	625 ± 33	(260)
	Mice		n. a. <sup>1</sup>	1,370 ± 150	(314)
	Rats		n. a. <sup>1</sup>	1,510 ± 250	(314)
	Rats		oral	4,290 (10% sol.)	*
	Rats		oral	2,600	(422a)
	Rabbits		n. a. <sup>1</sup>	1,900	(314)
	Rabbits		skin	0.5 ml/kg ( 500 mg/kg)	*
Sodium propionate	Mice		n. a. <sup>1</sup>	7,800	(314)
	Mice (DD-strain)		i. v.	1,380	(413)
			i. p.	2,250	(413)
			i. m.	3,200	(413)
			oral	5,100	(413)
	Rats		n. a. <sup>1</sup>	8,150	(314)
	Rabbit		n. a. <sup>1</sup>	6,750	(314)

<sup>1</sup>n. a. = not available

\*Private communication from unidentified source provided by FDA.



Table 1 (cont'd)

Substance	Animal and Strain	Number	Route	LD <sub>50</sub> (mg/kg body wt.)	Ref.
Calcium propionate	Mice (DD-strain)		i. v.	580	(413)
			i. p.	720	(413)
			i. m.	1, 020	(413)
			oral	3, 340	(413)
	Rats		oral	5, 160 (4, 170- 6, 380)	(546)
Thiodi- propionic acid	Rats		n. a. <sup>1</sup>	>2, 000	(566a)
Dilauryl thiodi- propionate	Rats		n. a. <sup>1</sup>	>2, 500	(422a)

Morphinized male albino rabbits weighing approximately 2 kg were used in eye irritation studies to observe the effect of chemical configuration on the edema-producing potency of acids and other compounds. The edema produced by propionate was quite similar to the amount produced with hydrochloric acid of the same molar concentration (196).

Acclimatization experiments were conducted in which the organisms were grown in nutrient broth containing 0.8% sodium propionate or sulfacetamide sodium. It was then sub-cultured for 4 days in broth with 1.25% of the separate drugs and the resultant cultures were used to inoculate broth containing various concentrations of the compounds. The results show that the organisms did not become appreciably resistant to sodium propionate. A test for synergism or antagonism was conducted using 2% sodium propionate + 2% sulfacetamide sodium. The results obtained with staphylococcus aureus suggest that antagonism does exist between the sodium propionate and sulfacetamide sodium preparations on local infections when propionate is applied topically to the eyes, ears, nose, vagina, vagina skin and other tissues (415).

Four-week-old white rats were used in a 4- to 5-week study of the toxicity of several food-preserving agents such as sodium propionate and calcium propionate. Weight gain was used as the indication of toxicity produced in animals fed diets containing 1 or 3% calcium or sodium propionate; neither additive affected the amount of weight gain (503).

To compare the utilization of energy, diets containing 24% acetate, propionate, or butyrate with glucose and starch were fed to weanling and adult female rats for periods of 5 or 20 days, respectively. Feeding of the propionate rations to the weanling animals was discontinued after 5 days due to high mortality and negative weight gain; the adult animals survived 20 days, but rough hair coats, reddened paws and extreme excitability were observed in all. Substitution of the propionate in the diet did not affect digestibility, metabolizability or energy measurements, but did interfere with normal metabolic function to the extent that it could not be fed to the younger rats and presented the aforementioned problems in the adults (231).

A series of studies were conducted to determine the fungicidal, fungistatic and antifermentative properties of sodium propionate. In the first study of the series, albino rats weighing approximately 200 g were given 100 mg/kg of propionate by hypodermic injection for a period of 5 weeks, while another group received 100 mg/kg by the same route of administration. Thereafter, there was a weekly dosage increase of 50 mg/kg until 350 mg/kg was reached. The results obtained show that there were no mortalities nor were there any noticeable variations in the weight of the treated animals at any of the dose levels when compared with the controls. After 4 weeks, there was a slight local loss of hair in the animals receiving the incremental doses, and in the 5th and 6th weeks, small subcutaneous abscessula protuberances

formed which may have been due to the large quantity of propionate administered. Analysis of the urine revealed that the volatile products of acid character underwent positive and negative variations of 3% in the control groups, whereas in the 2 medicated groups described, the same content increased by 11% and 16.5%, respectively. This finding led the authors to believe that part of the sodium propionate is eliminated through the urine in the form of volatile components without undergoing any preventive block within the organism. The second experiment of this series on the fungicidal activity of sodium propionate in vitro revealed that a .0125% to 2% solution of sodium propionate is one of the potent inhibitors of bacteria and fungi responsible for local and more serious infections. In the third series of experiments, apple juice, sweet filtrate of grape and orange juice were bottled and stored at temperatures between 20° and 30°. Either 0.2, 0.4, 0.6, or 0.8% by weight of sodium propionate was added to each group of 2 juices, and the samples were stored for approximately 10 months. The results show that 0.2, 0.4, and 0.6% concentrations would block fermentation for 5, 11, and 21 days, respectively, but would exert almost no action after that time (461).

Weanling pigs were administered propionic acid or calcium propionate, ( to give 0.8% propionate), in order to determine the effect on performance and bacterial flora. Treatments were applied for 4 weeks, and performance was also measured for a further period of 8 weeks after treatment had ceased. In a second study, a pig from each of 9 litters was

slaughtered for examination before treatments were applied in order to obtain bacterial baseline data. The remaining animals were placed on the treatment as described above for a period of 4 weeks, after which samples for bacterial examination were taken from the duodenum and jejunum. These studies showed that after 4 weeks, there was no significant difference in the growth rate of the pigs, and that bacteriologically, both substances tested were effective in controlling hemolytic *Escherichia coli* and reducing non-hemolytic *Escherichia coli* in the duodenum and jejunum (57).

A preliminary feeding trial of 3-weeks duration indicated that there was little difference in acceptability of several diets by weanling rats. Therefore, a long-term study was conducted in male and female rats to determine the chronic toxicity of bread additives. The findings showed that the baked bread product, (5% concentrations of propionate were added to bread ingredients before baking), fed ad libitum to rats as 3/4 of the diet for a one-year period, did not manifest any toxic effects. A follow-up study was conducted in which animals were fed a diet containing 75% sodium propionate for 32 weeks. It was found that sodium propionate at this high level caused a depression of growth during the first weeks of feeding, but did not influence mortality and produced no detectable effect on hemoglobin levels, organ weights, or histopathology of the tissues (412).

Mixed-strain male and female albino rats, weighing 50 to 80 g, were fed a diet containing 5 to 10% propionic acid to determine the

production of gastric lesions by these diets. One rat on the propionic acid diet died early in the experiment, and the remaining 4 were sacrificed after 110 days. In 3 or 4 cases it was found that umbilicate or warty lesions were evidenced in the forestomach, while the remaining one showed no gastric changes (237).

Male and female shorn lambs weighing 23 to 35 kg were fed a ration containing 10% sodium propionate twice daily for 50 days ad libitum. It was found that neither rate of body weight gain or feed required/unit gain was affected by incorporating 10% sodium propionate in the feed. At slaughter, it was found that the animals fed the propionate-containing diets were considerably higher in fat than the controls indicating that they converted digestible energy to weight gain considerably more than did the controls (85).

A study utilizing human skin and fetal lung cells and strain L cells was conducted to determine the effects of sodium salts and certain monocarboxylic acids on established cell lines. The results revealed that at a concentration of 10 mg%, sodium propionate was toxic to all cell lines tested, but had no significant effect at a 1 mg concentration (442).

Walter Reed-Carworth Farm rats in their first gestation were administered a total dose of 0.5 g thiopropionic acid during a 22-day diet period. On the 22nd day, the young were Caesarean delivered, and both horns of the uterus were excised and carefully

microscopically for resorption sights in order to determine if certain anti-oxidants influenced resorption in rats. The results show that, of 10 litters, there were 128 implantations, 119 were fetuses, and 7% were resorptions (575).

In vitro anti-tumor studies were carried out with 6 C3HED lymphosarcoma cells. Ehrlich ascites carcinoma and TA3 mammary carcinoma were injected intraperitoneally in C3H mice or Connaught mice weighing 20 to 22 g. The washed tumor cell suspension was added to 1 ml of Fink's solution containing 1.6 mg of propionic acid, and 0.5 ml of this suspension was administered intraperitoneally to mice. The results show that the survival time of the test animals was significantly longer than the control animals; after 90 days, 2 of the 10 animals medicated were alive (347).

The inhibition of tumor growth with antimetabolites of the hexose monophosphate pathway intermediates was studied in transplantable Yoshida ascite sarcoma cells in rats and fibrosarcoma cells in Swiss mice. Thiodipropionic acid was injected intraperitoneally 24 hours after the transplantation of the Yoshida sarcoma cells in rats, while the compound was injected subcutaneously near the solid fibrosarcoma in the mouse on the 6th day after transplantation. Thiodipropionic acid completely destroyed the sarcoma cells after 10 days of treatment, and the animals lived a normal life span without any trace of malignancy, while the corresponding control animals died within 10 days (571).

Male Wistar rats were kept at 25°C for 10 days and on the 11th day were given 1 g of tripropionin and 1.9 g of dextrose containing  $-1-C_{14}$ . The rats were then placed in a metabolism cage, and their respiratory  $CO_2$  was measured for 24 hours, after which the animals were sacrificed and head and carcass were analyzed for lipids. In a second study, animals were injected intraperitoneally with 1 ml of saline containing  $-1-C_{14}$  sodium propionate. They were placed at once in wire cages, either fed or fasted, were sacrificed after 24 hours, and the lipids were isolated and analyzed for  $C_{14}$ . The results showed that in the animals medicated with labeled sodium propionate by stomach tube, 75% of the  $C_{14}$  content appeared in the respiratory  $CO_2$  during the 24-hour experiment. In the animals which were administered labeled propionate intraperitoneally, very little of the  $C_{14}$  appeared in the body fat. A third study showed that the rat liver was able to convert large amounts of acetate into fatty acids but was unable to convert any of the carboxyl or propionate into long-chain fatty acids. In the study of the adipose tissue, it was evident from the position in which the propionates were labeled that propionate 2 or 3 carbons are far better precursors of long-chain fatty acids than is the carboxylcarbon (226).

A section of rat jejunum approximately 30 cm long was inverted to form a sack which was suspended in 15 to 50 ml of saline solution in order to study the transfer of short-chain fatty acids by the intestines. Transfer experiments, to measure changes in the mucosa,



and metabolism experiments to estimate the total amount of fatty acids disappearing during the course of the experiment were conducted using propionic acid. Concentrations of propionic acid added to the baths ranged from 5 to 80 mm. It was found that propionic acid was transferred against the concentration gradient, that high concentrations of propionic acid had an inhibitory effect on transfer of the fatty acid, and that the concentration necessary to cause this inhibition decreased with increasing length of the fatty acid chain (22).

Another study, by different authors using the same techniques as above but utilizing both jejunal and ileal sections, showed that the preferential movement for propionic acid was in the jejunum rather than the ileum (55).

For 2 weeks, twelve adult brown leghorn chickens were fed a diet containing approximately 2.5% propionate. Upon examination of the contents of gut sections, it was found that fatty acids were completely removed from the ingesta before it reached Meckel's diverticulum (35).

A study was conducted to determine the nature of the gastric mucosal entry by fatty acid and acetylsalicylic acid in vagally denervated pouches in the oxyntic gland area. Results obtained by 2 to 30 minute periods of irrigation of the denervated pouches with a solution containing 100 millimols of propionic acid revealed that short-chain fatty acids must be in fat soluble form before they

can diffuse through the mucosal barrier, and that once through the barrier, they cause damage to the mucosa (63).

The gastrocnemius muscle of the dog was isolated and transferred to a oxygenated blood circuit for perfusion. Isotopic sodium propionate was injected, and the collection of respiratory  $\text{CO}_2$  in sodium hydroxide absorbers was begun. The labeled propionate was injected into the bath at levels of 5, 10, 15, 20, and 25 millimols/l, and the results show that 9 to 13% of the respiratory  $\text{CO}_2$  was derived from the carboxyl carbon of propionate in the resting skeletal muscle preparation and only 3-5% in the repetitively contracting gastrocnemius. Evidence was obtained for the conversion of carboxyl-labeled propionate to carboxyl lactate by isolated skeletal muscle but not for its incorporation into muscle glycogen. When acetate and propionate were administered simultaneously to the resting muscle, the rate of conversion of the carboxyl of the propionate to  $\text{CO}_2$  was reduced to a much greater extent than that of the carboxyl carbon of acetate (207).

To study the deep depression of feed intake by injected metabolites, goats were fitted with ruminal fistulas and were infused intraruminally for 2 days with 2.5, 5, 10, and 20 millimols/minute of propionic acid. The results showed that feed intake was decreased, but there was no difference between the effect caused by sodium propionate or propionic acid when tested by a paired t-test. Sodium propionate ingestions caused an increase in oral water intake as the rates of sodium injection increased, whereas propionic acid injections caused reduced oral water intake with increasing rate of injections (393).

Holstein and Brown Swiss dairy cows were used to study sodium propionate's effect on milk and fat production, roughage consumption, blood sugar, and blood ketones. The results showed that feeding 1/2 pound of sodium propionate daily to these animals did not affect milk production, hay consumption, blood sugar levels, or weight gains, but there was a slight yet not significant decrease in butter fat. A supplemental study to determine the fatty acid content of rumen fluid was conducted in two fistulated animals fed propionate, and the results showed that there were higher than normal levels of all volatile fatty acids in the rumen for the first 3 hours after feeding. Calculations showed that all of the administered propionate had disappeared in 7 hours (458).

Propionic acid was administered intravenously by constant drip for 8 hours on three consecutive days to study appetite inhibition in dairy cattle by certain intermediate metabolites. Administration of the propionic acid resulted in a significant reduction in intake during 8-hour and 24-hour periods, but there was very little carry-over since eating began very soon after cessation of each injection period (72).

When 1.65 millicuries of radio-labeled propionate was infused through a teat cannula into the udder of lactating cows to study the synthesis of milk fat in the bovine mammary gland, the results showed that there was no synthesis of glycerol in the mammary gland caused by the infusion of propionate (215).

Twin Holstein cows and a Guernsey steer were fitted with fistula plugs and used in studies to determine the amounts of short-chain acids formed during rumen fermentation. The concentration of propionic acid in the rumen was determined at various times after feeding either an all-hay ration or one consisting largely of concentrates, and the results revealed that the concentrations of each acid first increased and then decreased after feeding. The data also revealed that the rate of disappearance of propionic acid in high-concentrate rations was 4.6%/hour, and 3.2%/hour in the hay ration (87).

Six female Heidehain pouch dogs with total antrum resections were used to study the stimulation of gastric acid secretion. The results show that propionate administered by venoclysis in a dose of 2.5 to 5.0  $\mu\text{m}/\text{kg}/\text{minute}$  had a pronounced stimulatory effect on the secretion of gastric acid from the pouches (435).

Rumen abomasal Crocker-Markowitz fistulas, oesophagostomy and venous catheterization were performed on 1- to 2-hour-old Clun-Forest sheep to obtain information about factors responsible for initiating and maintaining abomasal secretion. 250 millimols of propionic acid, neutralized to pH 5.5 with sodium hydroxide, were added to rumen contents at hourly intervals; the results show that propionic acid stimulated abomasal secretion (138).

To determine the biological activity of some acids, a cat was injected intravenously with a dose of  $1 \times 10^{-4}$  m/kg of sodium propionate. To study the in vivo and in vitro effects, a rat uterus was bathed with a  $10^{-6}$  solution of sodium propionate. The results showed that injected into the cat, sodium propionate caused hypertension and a pressure decrease (about 40 mm of mercury) for 1 minute. It was indicated that sodium propionate had no effect on the perfused rat uterus (451).

Merino wethers, 2 to 3 years of age, were infused for 60 minutes through the portal vein with solutions of  $C_{14}$ -labeled sodium propionate at the rate of 0.1 microcuries/minute/kg of body weight. Blood samples were drawn from the right jugular vein at termination of the infusion to determine the metabolism of propionic acid. At the end of the infusion, samples of liver and muscle were removed for glycogen assay. The results showed that the specific radioactivity of liver and muscle glycogen was low after the infusion of the labeled propionate. The  $C_{14}$ -1-propionate gave rise to labeling mainly in carbon 3 and 4 of glucose and carbon 1 of lactate, while  $C_{14}$ -2-propionate labeled mainly carbon 1, 2, 5, and 6 of glucose and carbon 1 and 2 of lactate (8).

Friesian cows fitted with large, permanent rumen fistulas were used in an infusion study of the effect of volatile fatty acid and lactic acid on the secretion of the component fatty acids in milk fat and on blood composition. In the first study, propionic acid

was infused into two separate cows at a rate of 1500 ml/day for one-week periods in order to study the fat content and fatty acid composition of milk. The results showed that during the infusion of propionic acid, palmitic acid was unaltered, but the yield of all other major fatty acids, with the possible exception of lauric acid, was decreased. In the second, third, and fourth series of experiments, propionic acid was infused at a rate of 1600 ml/day for 4 weeks to correlate blood composition as related to the changes observed in milk fat content. The results showed that propionic acid increased the concentration of glucose and decreased that of ketone body. Propionic acid caused only small increases in the volatile fatty acid content of the blood, indicating that the added acids were almost completely metabolized in the rumen epithelium or in the liver (334).

To demonstrate the utilization of propionate for hemesynthesis in human bone marrow cells, normal and megaloblastic bone marrow cells were compared in vitro with respect to the incorporation of C<sub>14</sub>-labeled propionate. In megaloblastic bone marrow cells obtained from patients with Vitamin B<sub>12</sub> deficiency, the incorporation of carbon-labeled propionate into heme was markedly reduced. The addition of coenzyme B<sub>12</sub> specific for a Vitamin B<sub>12</sub>-deficient state was found to be not directly related to morphologically identified megaloblastic changes (528).

The absorption of short-chain fatty acids was studied in a patient whose operation had left him with a fistula. The colon was profused with a solution containing acetic, propionic, butyric, isobutyric, valeric, isovaleric and caporic acids in equimolar amounts. The results show that absorption of short-chain fatty acids from a human's jejunum and colon depends on the chain length; the longer the chain, the faster the acids were absorbed (65).

To study the role of propionic acid as a precursor of methylmalonic acid in normal and Vitamin B<sub>12</sub>-deficient men, fasting subjects were injected with 10 to 15 microcuries of C<sub>14</sub>-2-propionate intravenously and/or microcuries of C<sub>14</sub>-2-methylmalonic acid. In addition, one of the Vitamin B<sub>12</sub>-deficient patients also received a 5 g oral dose of sodium propionate 30 minutes prior to the i.v. injection of the labeled propionate. The results of the loading study indicated that propionate is a precursor of urinary methylmalonic acid and, therefore, intracellular methylmalonyl CoA in both normal and Vitamin B<sub>12</sub>-deficient humans. The injected activity found in urinary methylmalonic acid was small, and the turnover studies indicate 20 to 50% is the daily turnover in normal patients and 20% in Vitamin B<sub>12</sub>-deficient patients. The fate of the remainder is unknown (67).

For 10 days, adult Wistar rats were fed ad libitum a diet composed of 25 g casein, 10 g fat, 51 g dextrose, 3 g salt mixture, 5 g cellulose, 6 g brewers yeast, 0.04 g cod liver oil concentration

and 0.01 g alpha tocopherol. Liver sections were taken to study the pattern alteration of hepatic carbohydrate metabolism by the addition of propionate as a substrate to the tissue bath. The results showed that propionate reduced the  $C_{14}O_2$  production to about 53% of that produced in control liver slices, while reducing the incorporation of the  $C_{14}$  into lipids to 24% of the control values (95).

Slices of mammary glands and livers of lactating Addis-Sloanaker rats were used to study propionate metabolism using 3, 5, 10, and 50 micromols/incubation vessels of  $C_{14}$  propionate. The experiments showed that, in the presence of glucose,  $C_{14}$ -1-propionate was extensively incorporated into fatty acids by mammary gland slices. The incorporation of  $C_{14}$ -2 and  $C_{14}$ -3-propionate into fatty acids was somewhat greater than that of  $C_{14}$ -1-propionate. The composition of the fatty acids from all labeled carbons was similar, but those labeled at the -2 and -3 carbon also contained approximately 20% more even carbon acids. The water soluble products formed from propionate  $C_{14}$  were determined in mammary glands, and it was found that the major labeled compounds were glutamic acid and glutamine, but when determined in the liver slices, the major products were  $CO_2$  and glucose (169).

The conversion of propionate to liver glycogen in the intact rat was studied using isotopic propionate. Variouslly labeled isotopic propionate + glucose preparations were fed to fasted rats, and after a suitable period of time, the livers were removed, the glycogen



isolated and hydrolyzed to glucose, and the distribution of isotopes in the glucose determined. The results revealed that the isotopes from the carboxyl group of propionates were found to give detectable labeling only in carbon atoms 3 and 4 of the glucose, while propionate-containing isotopes in the alpha or beta position gave rise to glucose labeled, predominantly and equally, in carbons 1, 2, 5, and 6 (517).

Female goats were employed to determine whether the administration of the individual volatile fatty acids, and combinations thereof, caused changes in the blood glucose and ketone body levels consistent with the generally-accepted mechanism of metabolism. Propionic acid was administered orally by a constant rate pump and passed into the rumen by way of the nostrils. Intravenous administration was made with a constantly-supervised drip device through a plastic tube inserted in the jugular vein. The data obtained from these studies demonstrated that propionic acid counteracts the formation of ketone bodies and causes a marked increase in blood glucose (161).

To study the effects of volatile fatty acids, including propionic acid, on serum insulin concentration, 4 ewes, 2 to 5 years of age, were administered a propionate solution intravenously (neutralized to pH 7 with sodium hydroxide) at a dose of 2.5 millimols/kg. Blood samples were drawn from the jugular vein 30 and 50 minutes before the injection and 0, 5, 10, 20, 30, 40, 60, and 90 minutes after

injection and were analyzed for free fatty acids, glucose, total ketone body, and insulin. The intravenous administration of propionate caused a slight increase in serum insulin, a marked hypoglycemia, and an immediate free fatty acid reduction which lasted for at least 90 minutes after the injection (223).

Presented in the same report was a study of propionate's effect on the changes in pancreatic activity at varying ages. Three sets of twin lambs were administered 2.5 millimols/kg of propionate intravenously either at 12 to 16, 42 to 46, 84 to 88, or 168 to 172 days of age. The administration of propionate produced an elevation in blood glucose and an initially slight hyperglycemia at all ages, but it was most evident in the lambs between 168 and 172 days of age. It was also found that the blood glucose concentrations returned to normal values sooner in the animals that were 12 to 16 days of age (223).

Fasted, adult phlorhizin-treated ewes weighing 130 to 150 pounds were used to study the effects of orally-administered short-chain fatty acids on certain blood and urine constituents. The animals were subjected to a 48-hour fasting period and twice-daily subcutaneous injections of 1g of phlorhizin dissolved in 5 ml of ethylene glycol were given. After this period, fatty acids were administered orally, and blood and urine specimens were collected for a 48-hour period. It was found that when propionic acid was administered in a dose of 1 g-mol/100 pounds of body weight, there was a significant increase

in blood glucose levels at 1, 3, 6, and 24 hours after treatment, with the greatest increase seen at the end of the 3rd hour. The blood ketone level was significantly below the controls, and the urine showed a marked increase in urinary glucose excretion during the 48-hour period after treatment (118).

A chronic surgical preparation permitting local enrichment of the pancreatic arterial blood was used as part of an experiment designed to determine if propionate played an important physiological role in regulating insulin secretion in sheep. When infused at a rate of 0.8 millimols/minute into the jugular vein, propionate produced a rapid and prolonged insulin elevation from 50 to 165  $\mu$ /ml. The glucose levels did not change in any of the trials involving propionate (222).

Seven dairy calves, 4 to 7 months of age, were administered 3 ml of P-32-tagged erythrocytes which were injected into the coeliac artery and collected from the gastro-splenic vein. The difference in the amount of fatty acids in blood taken from the jugular vein and that taken from the splenic vein was taken as the measure of absorption. Based on the analysis of serial blood samples obtained, it was calculated that 25 g of propionic acid would be absorbed for each pound of dry feed consumed/animal during a 24-hour period (58).

Mixed white-faced lambs, Holstein calves 5 to 90 days old, and mature dogs and horses were used to study the effect of volatile fatty acids on plasma glucose concentrations. The acids, in 2.5

normal solutions neutralized with sodium hydroxide to pH 7.2, were administered intravenously in doses of 2.5 mm/kg of body weight and caused a small plasma glucose increase and a remission of hypoglycemia convulsions in lambs, but did not affect the plasma glucose levels in other species. The small number of animals and the complications involved in administering the compounds made it difficult for the investigators to assess the value of this particular study (275).

Lipid solubility, permeability and hemolytic action of the saturated fatty acids were studied in human and dog red blood cells. The results show that propionic acid was the second most effective of the fatty acids in penetrating the red blood corpuscles, thus producing hemolysis, and that increasing the osmotic concentration of the outside fluid retards hemolysis. Human corpuscles were found to be more resistant to hemolysis by fatty acids than dog corpuscles; the resistance was apparently due to the greater buffer content of washed human corpuscles resulting in the neutralization of a greater amount of acid (32).

In an experiment studying the influence of fatty acid ions on the electro-encephalogram patterns of adult male albino rabbits weighing approximately 2 kg were injected intravenously with a propionate solution at 4 millimols/kg of body weight which immediately induced changes in the EEG patterns from a resting type to one of sleep. This sleep pattern was similar to the EEG pattern observed

when the animals were under deep sodium penta-barbitol anesthesia (372).

The effects of metabolically-important substances on the rabbit auricle were determined by addition of substances to a bath in which the auricles were suspended. Propionates introduced to the bath showed a tendency to induce arrhythmias which were occasionally completely irregular but usually were of the pulus alternas type (amplitude of contractions with each beat regularly alternating between two values). Rapid recovery from these arrhythmias was observed when the propionate was removed from the bath (368).

## TABLE OF CONTENTS

<u>SUBJECT</u>	<u>PAGE</u>
Summary	1
Chemical Information	5
Acute Toxicity	21
Short-term Studies	23
Long-term Studies	26
Special Studies	28
<u>Bibliography</u>	
Propionic Acid	B-1
Sodium Propionate	B-34
Calcium Propionate	B-41
Thiodipropionic Acid	B-48
Dilauryl Thiodipropionic Acid	B-50

## PROPIONIC ACID

- (1) Aas, M., and J. Bremer. 1969. Activation of propionate in rat kidney. Fed. Eur. Biochem. Soc. Lett. 5(1): 57-59.
- (2) Abe, T., and H. Tsuyuki. 1969. Studies on organic acids in "shottsuru" (fish sauce). Journal of Food Science and Technology 16(12): 560-563.
- (3) Aiso, K. 1967. Toxicity and safety of food additives. Kagaku Kogyo 18(2): 1-5.
- (4) Albonico, F., et al. 1964. Ripening of cheese. Riv. Latte. 20(3): 1-38.
- (5) Allen, G.R., and M.J. Saxby. 1968. Gas chromatography of isomeric fatty acid methyl esters. J. Chromatogr. 37(2): 312-314.
- (6) Al-Shabibi, M.M.A., et al. 1964. Effect of added fatty acids on the flavor of milk. J. Dairy Sci. 47(3): 295-296.
- (7) Annison, A. F., D.B. Lindsay. 1962. Measurement of entry rates of propionate and of butyrate in sheep. Biochem. J. 85: 474-479.
- \* (8) Annison, A.F., et al. 1963. The metabolism of acetic acid, propionic acid and butyric acid in sheep. Biochem. J. 88: 248-252
- (9) Anon. 1960. Effect of propionic acid on the secretion of the abomasum in sheep. VI. Izvest. Inst. Sravnitelna. Patol. Domashnite Zhivotni, Bulgar. Akad. Nauk. 8: 131-147.
- (10) Anon. 1963. Evaluation of the toxicity of a number of antimicrobials and antioxidants. Food Cosmet. Toxic. 1: 37-39.
- (11) Antonini, J., and J. Adda. 1969. Estimation by thin-layer chromatograph of propionic and butyric acids in Comte and Emmental cheese. Annales de Technologie Agricole 18(2): 139-142.
- (12) Arai, S., et al. 1966. Flavor components in soybean. III. Volatile fatty acids and volatile amines. Agr. Biol. Chem. 39(9): 863-868.
- (13) Armstrong, D.G., and K.L. Blaxter. 1957. The utilization of acetic, propionic and butyric acids by fattening sheep. Brit. Jour. Nutrition 11(4): 413-425.

- (14) Ash, R.W., et al. 1959. Effect of short-chain fatty acids on blood glucose in sheep. *Biochem. J.* 71: 9-10.
- (15) Ash, R.W., et al. 1964. The effect of short-chain fatty acids on blood glucose concentration in sheep. *Biochem. J.* 90(2): 353-360.
- (16) Attaway, J.A., et al. 1964. Identification of alcohols and volatile organic acids from natural orange essence. *J. Agr. Food Chem.* 12(2): 118-121.
- (17) Bacci, V., and G. Mannelli. 1968. Minor components of aqueous alcoholic liquids. IV. Volatile acids in Italian wines. *Atti. Congr. Qual.*, 6th, 1967: 433-442.
- (18) Bach, D. 1932. Desinfektionsversuche. *Bull. Sci. pharmacol.* 39(34): 499-504.
- (19) Baglioni, A. 1937. Propionic, toxicity; experimental study. *Fisiol. e med.* 8: 347-354.
- (20) Baile, C.A., and C.L. McLaughlin. 1970. Feed intake of goats during volatile fatty acid injections into four gastric areas. *J. Dairy Sci.* 53(8): 1058-1063.
- (21) Bandelin, F.J. 1958. The effect of pH on the efficiency of various mold-inhibiting compounds. *J. Am. Pharm. Assoc.* 47: 691-694.
- (22) Barry, R.J.C., and D.H. Smyth. 1960. Transfer of short-chain fatty acids by the intestine. *J. Physiol.* 152: 48-66.
- (23) Bässler, K.H. 1959. Comprehensive reviews. Metabolism and metabolic action of propionic acid in view of its application as a food preservative. *Z. Lebensm.-Untersuch. u. -Forsch* 110: 28-42.
- (24) Bates, R.C., and R.E. Hurlbert. 1970. The effect of acetate on *Euglena gracilis* var. *bacillaris* as a function of environmental conditions. *J. Protozool.* 17(1): 134-138.
- (25) Beacham, L.M. 1946. Decomposition in canned oysters and clams. *J. Assoc. Official Agr. Chem.* 29: 89-99.
- (26) Becker, E. 1954. Paper chromatographic detection of water-soluble organic acids in foods. *Z. Lebensm.-Untersuch. u. -Forsch.* 98: 249-257.



- (27) Bekirski, D.M. 1969. Amino acids and organic acids in Armenian peaches. *Biologicheskii Zhurnal Armenii* 22(10): 116.
- (28) Belehraddek, J., and F. Schwarz. 1928. Toxic action of acids of the formic series in relation to adaptation of the organism. *Bull. Soc. Chim. Biol.* 10: 909-919.
- (29) Bellucci, G., and B. Grigatti. 1970. Separation and determination of some organic acids formed during the spontaneous aerobic fermentation of tomatoes. *Industria Conserve* 45(1): 11-14.
- (30) Black, A. L., et al. 1966. Pyruvate and propionate metabolism in lactating cows. Effect of butyrate on pyruvate mechanism. *J. Biol. Chem.* 241(22): 5233-5237.
- (31) Blinc, M., et al. 1956. Propionic acid bacteria as inhibitors in the prevention of ropiness in bread. *Brot. u. Gebäck* 10: 66-68.
- \* (32) Bodansky, M. 1928. Lipoidlöslichkeit, Permeabilität und hämolytische Wirkung von gesättigten Fettsäuren. *Journ. biol. Chemistry* 79: 241-250.
- (33) Boehmer, T. 1968. Formation of propionylcarnitine in isolated rat liver mitochondria. *Biochim. Biophys. Acta* 1968: 164(3): 487-497.
- (34) Bogdonoff, P.D., et al. 1965. Composition of feed for ducks. French Patent 1 383 733.
- \* \* (35) Bolton, W., and W.A. Dewar. 1965. The digestibility of acetic, propionic and butyric acids by the fowl. *Brit. Poultry Sci.* 6(2): 103-105.
- (36) Bomar, M. 1962. Mode of action of fungitoxic compounds on microorganisms. II. Synergism of the bactericidal effect of certain chemical preservatives and low temperatures. *Folia Microbiol.* 7: 298-305.
- (37) Bottazzi, V. 1955. Paper chromatography of the volatile fatty acids from C<sub>2</sub> to C<sub>8</sub> found in products used in cheese factories. *Lab. sci.* 3: 166-172.
- (38) Bové, J., and R. Raveux. 1957. Separation and determination of carboxylic acids from C<sub>1</sub> to C<sub>6</sub> by partition chromatography on a silica column. II. Proposed new technique. *Bull. soc. chim. France* 1957: 376-381.

- (39) British Patent 754 028. 1956. Erythromycin production.
- (40) Broadbent, G. A. , and J. J. Huitson. 1968. Antifungal treatment of cereal grains. French Patent 1 538 839
- (41) Brown, F. , and L. P. Hall. 1950. The separation of carboxylate ions on the paper chromatogram. Nature 166: 66-67.
- (42) Brown, G. W. , Jr. , et al. 1956. The oxidative metabolic pattern of mouse hepatoma C<sub>954</sub> as studied with C<sup>14</sup>-labeled acetates, propionate octanoate, and glucose. Cancer Research 16: 509-519.
- (43) Buchanan, J. M. , et al. 1943. The role of carboxyl-labeled acetic, propionic, and butyric acids in liver glycogen formation. Jour. Biol. Chem. 150(2): 413-425.
- (44) Califano, L. 1937. "Über Fettsäureoxydation bei der Fettleber. Biochem. Z. 289: 354-364.
- (45) Canic, V. D. , and N. Perisic-Janjic. 1970. Separation of organic acids by thin-layer chromatography on starch. Tehnika 25(2): 330-332.
- (46) Cannata, J. J. B. , et al. 1965. Metabolism of propionic acid in animal tissues. XII. Properties of mammalian methylmalonyl coenzyme A mutase. J. Biol. Chem. 240(8): 3249-3257.
- (47) Cantoni, C. , et al. 1965. The action of the Micrococcaceae on pork fat: C<sub>1-5</sub> volatile fatty acids. Ind. Conserve 40(3): 205-207.
- (48) Cantoni, C. , et al. 1969. Putrefaction of sausages. Archivio Veterinario Italiano 20(4): 245-264.
- (49) Cantoni, C. , et al. 1969. Souring of uncooked ham. Archivio Veterinario 20(5): 355-370.
- (50) Carelli, V. , et al. 1955. Sorption chromatography of polar substances on polyamides. Nature 176(4471): 70-71.
- (51) Cats, H. , and H. Onrust. 1958. Identifying food preservatives by paper chromatography. Chem. Weekblad 54: 456-459.
- (52) Chi, Y. F. , and Y. T. Wong. 1934. Chemische Untersuchung der Wurzel von Umbellifera Peucedanum decursivum, Maxim. J. Chin. chem. Soc. 2: 329-336.

- (53) Clark, E.P., and F. Hillig. 1938. The Dyer method for the identification and determination of volatile fatty acids. J. Assoc. Official Agr. Chem. 21: 684-688.
- (54) Clarke, B.J., et al. 1962. Trace volatile constituents of beer. IV. Volatile acids. J. Inst. Brewing 68: 179-187.
- \* (55) Clarkson, T.W., et al. 1961. Transport of univalent anions by isolated small intestine of the rat. Am. J. Physiol. 200: 781-788.
- (56) Cohen, P.P. 1937. Untersuchungen über Ketogenese. J. biol. Chemistry 119: 333-346.
- \* (57) Cole, D.J., et al. 1968. The effect on performance and bacterial flora of lactic acid, propionic acid, calcium propionate and calcium acrylate in the drinking water of weaned pigs. Vet. Rec. 83: 459-464.
- \* (58) Conrad, H.R., et al. 1958. Estimating gastrosplenic blood flow and volatile fatty acid absorption from the forestomachs of calves. Jour. Dairy Sci. 41(8): 1094-1099.
- (59) Cooper, E.A., and A.E. Goddard. 1954. Solvent influences in disinfection. Chemistry & Industry 1954: 695-696.
- (60) Corrao, A. 1957. Researches on the soluble volatile acids of olive oil. Olii minerali, grassi e saponi, colori e vernici 34: 536-538.
- (61) Corse, D.A., and E.J. Murray. 1970. Propionate utilization by pregnant, lactating, and spontaneously ketotic dairy cows. J. Dairy Sci. 53(6): 740-746.
- (62) Daus, L., et al. 1952. Propionic acid metabolism in mouse-liver slices. J. Biol. Chem. 196: 77-87.
- \* (63) Davenport, H.W. 1964. Gastric mucosal injury by fatty and acetylsalicylic acids. Gastroenterology 46: 245-253.
- (64) Davies, C.W., and B.D.R. Owen. 1956. Adsorption of organic acid molecules and their chromatographic separation on ion-exchange resins. J. Chem Soc. 1956: 1681-1685.

- (65) Dawson, A. M., et al. 1964. Absorption of short-chain fatty acids in man. Proc. Soc. Exptl. Biol. Med. 117(1): 97-100.
- (66) Declercq, B., et al. 1969. Gas chromatographic determination of volatile acids in effluent waters. Int. Symp. Chromatogr. Electrophoresis, 5th 1968: 130-137.
- (67) DeGrazia, J., et al. 1969. The role of propionic acid as a precursor of methylmalonic acid in normal and vitamin B<sub>12</sub>-deficient man. J. Lab. Clin. Med. 73(6): 917-923.
- (68) Deuticke, H. J. 1924. Über die Bedeutung von Ionen für die Muskelfunktion. VII. Die Wirkung verschiedener organischer Anionen auf den Lactacidogenwechsel. Ztschr. f. physiol. Ch. 141: 196-224.
- (69) Dickson, W. H., and M. J. Wilson. 1928. Weitere Beobachtungen über die Motilität des menschlichen Magens. Journ. Pharmacol. exptl. Therapeutics 34: 65-72.
- (70) Diemair, W., and E. Schams. 1960. Gas chromatography in food analysis. I. Determination of the lower, volatile fatty acids in foods. Z. Lebensm.-Untersuch. u. -Forsch. 112: 457-463.
- (71) Dilanyan, Z. K., and G. F. Genich. 1970. Accumulation of flavour and aromatic compounds in Ukrainian cheese. Molochnaya Promyshlennost' 31(10): 15-18.
- \* (72) Dowden, D. R., and D. R. Jacobson. 1960. Inhibition of appetite in dairy cattle by certain intermediate metabolites. Nature 188: 148-149.
- (73) Dowden, B. F., and H. J. Bennett. 1965. Toxicity of selected chemicals to certain animals. J. Water Pollution Control Federation 37(9): 1308-1316.
- (74) Drews, E. 1953. Detection of organic acids and preservatives by means of paper chromatography. Getreide u. Mehl 3: 85-88, 90-91.
- (75) Dubos, R. J. 1950. The effect of organic acids on mammalian tubercle bacilli. J. Exptl. Med. 92: 319-332.
- (76) Dudzinski, A. E. 1967. A spray sequence specific for the detection of free fatty acids. J. Chromatogr. 31(2): 560.

- (77) Dumazert, C., and C. Ghiglione. 1959. Chromatography by steam distillation. *Bull. soc. chim. France* 1959: 615-617.
- (78) Duncan, R.E.B., and J.W. Porteous, 1953. Identification and determination of the lower straight-chain fatty acids by paper partition chromatography. *Analyst* 78: 641-646.
- (79) Dutton, P.L., and W.C. Evans. 1970. Inhibition of aromatic photo-metabolism in *Rhodopseudomonas palustris* by fatty acids. *Arch. Biochem. Biophys.* 136(1): 228-232.
- (80) Efthymiou, E. 1967. Major free fatty acids of Feta cheese. *J. Dairy Sci.* 50(1): 20-24.
- (81) Egan, A.R. 1966. Nutritional status and intake regulation in sheep. V. Effects of intraruminal infusions of volatile fatty acids on voluntary intake of roughage by sheep. *Australian J. Agr. Res.* 17(5): 741-755.
- (82) Ege, R. 1924. Untersuchungen über die Durchdringungsfähigkeit der Anionen bei Berührung mit der Membran der roten Körperchen. *C. r. soc. de biologie* 91: 779-781.
- (83) Eichholtz, V.F., and E. Brandes. 1937. Iron catalysis. *Arch. Exptl. Path. Pharmacol.* 187: 215-220.
- (84) Elema, J. 1966. Comparison of analytical methods for the determination of volatile fatty acids, particularly propionic acid. *Z. Tierphysiol. Tierernaehr. Futtermittelk.* 21(6): 345-350.
- \* \* (85) Elliot, J.M., et al. 1965. Effect of acetate and propionate on the utilization of energy by growing-fattening lambs. *J. Nutr.* 87(2): 233-238.
- (86) Elsdon, S.R. 1946. Application of the silica gel partition chromatogram to the estimation of volatile fatty acids. *Biochem. J.* 40: 252-256.
- \* (87) Emery, R.S., et al. 1956. The amounts of short chain acids formed during rumen fermentation. *Jour. Animal Sci.* 15(3): 854-862.
- (88) English Patent 455 846 vom 23/3 1936. 1936. Verhinderung von Schimmelbildung auf Brot.

- (89) Erwin, E.S., et al. 1961. Volatile fatty acid analyses of blood and rumen fluid by gas chromatography. *J. Dairy Sc.* 44: 1768-1771.
- (90) Eusebio, A.N., et al. 1959. Effect on rumen volatile fatty acids and rumen microbial dissimilation by glucose-C<sup>14</sup> of corn meal when fed exclusively and in combination with hay or certain additives. *Jour. Dairy Sci.* 42(4): 692-697.
- (91) Fabricus, N.E. 1940. The manufacture and use of butter culture. *Dairy Inds.* 5: 159-160.
- (92) Fairbairn, D., and R.P. Harpur. 1951. Distillation, extraction, and chromatographic separation of the C<sub>2</sub>-C<sub>8</sub> fatty acids. *Can. J. Chem.* 29: 633-641.
- (93) Falchuk, M.Z., and G.W.G. Sharp. 1968. The role of the tricarboxylic acid cycle in the stimulation of sodium transport by aldosterone. *Biochim. Biophys. Acta* 153(3): 706-714.
- (94) Favarger, P., and J. Gerlach. 1966. Research on metabolism of odd-numbered fatty acids and the synthesis of fatty acids from propionic acid. *Helv. Chem. Acta* 49: 506-509.
- \*(95) Felts, J.M., et al. 1958. Alteration of the pattern of hepatic carbohydrate metabolism by propionate and butyrate. *Arch. Biochem. Biophys.* 75: 361-367.
- (96) Fernholz, H., and H. Neu. 1971. Process for preserving bakery products with mixed propionic acid anhydrides. West German Patent Application 1 793 221.
- (97) Ferretti, A., et al. 1970. Nonenzymic browning in a lactose-casein model system. *J. Agr. Food Chem.* 18(1): 13-18.
- (98) Ferretti, A., and V.P. Flanagan. 1971. Nonenzymatic browning in edible spray-dried whey. Identification of some volatile components. *Journal of Dairy Science* 54(12): 1769-1771.
- (99) Fischer, M.H., and W.J. Suer. 1934. Der Einfluss verschiedenartiger anorganischer und organischer Säuren und Oxysäuren auf die Quellung von Eiweiskörpern. *Arch. Pathology* 17: 615-619.
- (100) Flatt, W.P., et al. 1969. Influence of pregnancy and ration composition on energy utilization by dairy cows. *Eur. Ass. Anim. Prod. Publ.* 1967: 12: 123-138.

- (101) Flavin, M., et al. 1955. Metabolism of propionic acid in animal tissues. *Nature* 176(4487): 823-826.
- (102) Flavin, M., and S. Ochoa. 1955. Propionic acid metabolism in animal tissues. *Congr. intern. biochim., Resumes communes.*, 3 Congr. Brussels, 1955: 51.
- (103) Flavin, M., et al. 1956. Metabolism of propionic acid in animal tissues. *Biochem. Problems Lipids, Proc. Intern. Conf.* 2nd, Ghent 1955: 208-211.
- (104) Flosner, O. 1931. Beitrage zur Biochemie menschlicher Korperflussigkeiten. II. Untersuchungen über Ovarialzystomflüssigkeiten. *Ztschr. Biol.* 91: 291.
- (105) Fonda, M. L., and B. M. Anderson. 1968. D-Amino acid oxidase. II. Studies of substrate-competitive inhibitors. *J. Biol. Chem.* 243(8): 1931-1935.
- (105a) Food Chemicals Codex. 1972. 2nd. Ed. National Academy of Sciences. Washington, D. C. pp. 151-152, 676-677, 761, 1246-1247.
- (106) Fourton, A. 1928. Existiert die Antirrhinsäure in der Digitalis? *Bull. Sciences pharmacol.* 35: 689-691.
- (107) Frank, M. and E. Hess. 1941. Studies on salt fish. V. Studies on *Sporendonema epizoum* from "Dun" salt fish. *Jour. Fish. Res. Bd. Canada* 5(3): 276-286.
- (108) Freixo, J. 1958. Characteristics of cod low in salt. *Conservas de Peixe* 13: No. 152: 18.
- (109) French Patent 1 587 601. 1970. Compositions for fish conservation.
- (110) Furchgott, R. F., and E. Schorr. 1946. Sources of energy for intestinal smooth muscle contraction. *Proc. Soc. Exptl. Biol. Med.* 61: 280-286.
- (111) Garcia, P. T., and L. Cuerpo. 1966. A study of alterations in the composition of volatile fatty acids derived from ruminal liquor from the time of extraction to the time of analysis. *Rev. Invest. Agropecuar. Ser. I*: 3(4): 69-80.
- (112) Georges, A. 1954. Bacteriostatic and fungistatic activities of propionic and caprylic acids as a function of pH. *Excerpta Med., Sect. XIII*: 8: 80.

- (113) Gerisma, S.Y. 1970. Gelatinizing temperatures of starch, as affected by polyhydric and monohydric alcohols, phenols, carboxylic acids, and some other additives. *Staerke*, 22(1): 3-9.
- (114) Gerstley, J.R., et al. 1928. The influence of feeding on certain acids in the feces of infants. A comparison of the effects of breast milk and modified cow's milk on the excretion of volatile acids. *Amer. Jour. Dis. Children* 35(4): 580-589.
- (115) Ghorban, K.Z., et al. 1966. Concentrations of volatile fatty acids and lactic acid in the rumen as influenced by diet and post-feeding time. *J. Dairy Sci.* 49(12): 1515-1518.
- (116) Gianturco, V.M., et al. 1966. Volatile constituents of coffee. *Nature* 210(5043): 1358.
- (117) Goepfert, J.M., and R. Hicks. 1969. Effect of volatile fatty acids on *Salmonella typhimurium*. *J. Bacteriol.* 97(2): 956-958.
- \* (118) Goetsch, G.D., and W.R. Pritchard. 1958. Effects of oral administration of short-chain fatty acids on certain blood and urine constituents of fasted, phlorhizin-treated ewes. *Amer. Jour. Vet. Res.* 19(72): 637-641.
- (119) Gogiya, V.T. 1965. Chromatographic study of some components of tea essential oil. *Chai, Kul'tura i Prozv.*, Tbilisi, Sb. 1965(1): 56-67.
- (120) Gogiya, V.T., and L.N. Ivanova. 1968. Composition of pink geranium essential oil. *Prikl. Biokhim. Mikrobiol.* 4(6): 733-738.
- (121) Gonthier, B., and J. Mocotte. 1970. Biocidal preservatives for fish. German Patent 1 921 670.
- (122) Gordillo, A.L., and A.L. Montes. 1960. Chromatography of fatty acids; semimicro method for determination of C<sub>1</sub>-C<sub>10</sub> acids. *Anales soc. cient. arg.* 170: No. 3-4: 53-67.
- (123) Grafflin, A.L., and D.E. Green. 1948. Cyclophorase system. II. Complete oxidation of fatty acids. *J. Biol. Chem.* 176: 95-115.
- (123a) Graveland, A. 1972. A gas Chromatographic determination of propionic, sorbic and benzoic acid in rye bread and margarine. *J. Assoc. Off. Anal. Chem.* 55(5): 1024.
- (124) Green, J.W. 1949. The relative rate of penetration of the lower saturated monocarboxylic acids into mammalian erythrocytes. *J. Cellular Comp. Physiol.* 33: 247-266.



- (125) Grimmer, W, and E. Fink, 1934. Beitrage zur Biochemie der Mikroorganismen. IV. Zur Kenntnis der Acidopropionolactone. Milchwirtschaftl. Forschg. 16: 277-288.
- (126) Grosjean, M.H., and A. Fouassin. 1967. Gas-phase chromatographic determination of propionic acid in foods. Rev. Ferment. Ind. Aliment. 22(6): 211-213.
- (127) Guggenheim, B., et al. 1965. Effect of volatile acids in saliva on the aerobic metabolism of bacteria in mouth. Pathol. Microbiol. 28: 77-83.
- (128) Gulați, B.C. 1969. Essential oils of wild ginger of Kumaon. Perfum. Essent.Oil Rec. 60(78): 257-258.
- (129) Gunthier, B., and J. Mocotte, 1971. Fish preservation. United States Patent 3 600 198.
- (130) Hamid, H.A., and R.L. Saffle, 1965. Isolation and identification of the volatile fatty acids present in hickory sawdust smoke. J. Food Sci. 30(4): 697-701.
- (131) Hamid, H.A., and R.L. Saffle, 1966. Penetration depth of volatile fatty acids into fresh ham. Food Technol. 20(2): 126-127.
- (132) Hansen, J.D. and M.D. Appleman. 1955. Effect of sorbic, propionic, and caproic acids on growth of certain clostridia. Food Res. 20: 92-96.
- (133) Hawke, J.C. 1957. The fatty acids of butterfat and the volatile acids formed on oxidation. Jour. Dairy Res. 24(3): 366-371.
- (134) Henning, S.J., and F.J.R. Hird. 1970. Concentration and metabolism of volatile fatty acids in the fermentative organs of two species of kangaroo and the guinea pig. Brit. J. Nutr. 24(1): 145-155.
- (135) Hernandez-Gutierrez, F. 1956. A study of preservatives and antiseptics used for flour and its derivatives. Anales bromatol. 8: 39-72.
- (136) Hibbs, J.W., et.al. 1956. A high roughage system for raising calves based on early development of rumen function. VI. Influence of hay to grain ratio on calf performance, rumen development, and certain blood changes. J. Dairy Sci. 39: 171-179.

- (137) Hidaka, T., et al., 1965. Behavior of organic acids in hot air drying of sweet potato starch. Denpun Kogyo Gakkaishi 12(3): 95-100.
- (138) Hill, K.J. 1960. Abomasal secretion in the sheep. J. Physiol. 154: 115-132.
- (139) Hillig, F., 1944. Report on the determination of volatile acids in fish and fish products. J. Assoc. Official Agr. Chem. 27: 237-240.
- (140) Hillig, F. 1945. Report on [the determination of] acid in eggs. J. Assoc. Official Agr. Chem. 28: 217-221.
- (141) Hillig, F., et al. 1949. Water-insoluble fatty acids and butyric acid in cream and butter. J. Assoc. Official Agr. Chemists 32: 731-745.
- (142) Hoffsten, R., and M. Kellner. 1971. Processing of chemically preserved wet maize in the distillery. Branntweinwirtschaft 111(1): 1-4.
- (143) Hollenback, C.M., and L.J. Marinelli. 1963. Some chemical constituents of smoke flavoring. Proc. Res. Conf. Advisory Council Res. Am. Meat Inst. Found. Univ. Chicago 15: 67-76.
- (144) Holter, J.B., et al. 1970. Heat increments of steam-volatile fatty acids infused separately, and in a mixture, into fasting cows. J. Dairy Sci. 53(9): 1241-1247.
- (145) Hori, S., and A. Tanabe, 1956. Quantitative estimation of straight-chain volatile fatty acids by column partition chromatography. Okayama Daigaku Nogakubu Gakujutsu Hokoku No. 9: 47-54.
- (146) Huitson, J.J., and P.W.J. Watts. 1968. Prevention of mold in animal feeds. French Patent 1 549 021

- (147) Hunter, I.R., et al. 1960. Gas-chromatographic separation of volatile organic acids in the presence of water. *Anal. Chem.* 32L 682-684.
- (148) Huntington, G.I., and O. Rahn. 1945. Acidity controlling antiseptics by weak acids. *Jour. Bact.* 50(6): 655-659.
- (149) Ichikawa, Y., and Y. Kitamoto. 1967. The antimicrobial action of propionate and related substances. II. The inhibitory properties of propionate and related substances on the growth of *Escherichia coli*. *Nippon Nogei Kagaku Kaishi* 41(5): 171-177.
- (150) Ihne, H. 1970. Chemical or physical preservation of cereals? *Deutsche Muller-Zeitung* 68(4): 80-81.
- (151) Imai, S., and K. Suzuki. 1969. Components of shoyu mash brewed at different temperatures in the early stage. *Seasoning Science* 16(5): 1-5.
- (152) Inagaki, C. 1943. Vitamin C in citrus fruits. I. The comparison of the dried powder of natsumikan juice and synthetic L-ascorbic acid in utilization by animal experiments. *J. Agr. Chem. Soc. Japan* 19: 163-174.
- (153) Inoue, Y., and M. Noda. 1950. Paper partition chromatography of aliphatic carboxylic acids by means of hydroxamic acid method. *J. Agr. Chem. Soc. Japan* 23: 368.
- (154) Inouye, S. 1925. Die Kennzeichnung der biologischen Wirkung von Saure auf das Froschherz. *Acta scholae med. Kioto* 7: 227-240.
- (155) Ishigaki, R., and I. Nagase. 1964. Semichemical soy sauce. III. Quantitative transition of amino acids in the semichemical processing of soy sauce. *Nippon Jozo Kyokai Zasshi* 59(2): 161-162.
- (156) Ivanov, N., and L. Stancheva. 1968. Effect of lower fatty acids on liver secretions in sheep. *C.R. Acad. Sci. Agr. Bulg.* 1968: 1(2): 173-177.
- (157) Iwata, N., et al. 1965. Rancid flavor of edible oils. I. Rancid flavor components of soybean oil. *Yukagaku* 14(5): 241-246.
- (158) Jackel, S.S. 1969. Fermentation flavours of white bread. *Abstracts of Papers, American Chemical Society* 158: MICR 41.

- (159) Jackson, R.B. 1966. Gas-liquid chromatography of volatile fatty acids from formic to valeric acid. II. The instability of silicone oil-fatty acid stationary phases. *J. Chromatog.* 22(2): 251-260.
- (160) Jacobson, D.R., et al. 1966. Appetite regulation in the ruminant. *Proc. Int. Congr. Nutr.*, 7th 1966: 2: 63-67.
- \* (161) Johnson, R.B. 1955. The effects of the administration of acetic, propionic, and n-butyric acids upon the blood glucose and ketone body levels of goats. *Cornell Vet.* 45(3): 273-289.
- (162) Johnson, R.R., et al. 1958. Studies on the in vitro rumen procedure: Improved inoculum preparation and the effects of volatile fatty acids on cellulose digestion. *Jour. Animal Sci.* 17(3): 841-850.
- (162a) Joint FAO/WHO Expert Committee on Food Additives 1966. Toxicological evaluation of some antimicrobials, antioxidants, emulsifiers, stabilizers, flour-treatment agents acids and bases. 9th Report. Report Series No. 40A, B, C, pp. 56-57.
- (163) Jones, D.D., and R.N. Greenshields. 1969. Volatile constituents of vinegar. I. Survey of some commercially available malt vinegars. *J. Inst. Brew.*, London 75(5): 457-463.
- (164) Jones, D.D., and R.N. Greenshields. 1970. Volatile constituents of vinegar. II. Formation of volatiles in a commercial malt vinegar process. *Journal of the Institute of Brewing* 76(1): 55-60.
- (165) Kaziro, Y., et al. 1964. The metabolism of propionic acid. *Advances Enzym* 26: 283-378.
- (166) Kami, T., et al. 1969. A study on the flavour components of meat and meat products. II. Low-boiling compounds in barrow's subcutaneous fat depot and fresh lean pork. *Journal of the Faculty of Fisheries and Animal Husbandry, Hiroshima University* 8(2): 105-117.
- (167) Kananadze, T.A. 1963. Separation of fatty acids by partition chromatography. *Vinodelie i Vinogradarstvo SSSR* 23(4): 8-10.
- (168) Kanno, S., et al. 1969. Analysis of food additives. XVIII. Determination of propionic acid in bread by liquid column chromatography using a thermal detector. *Shokuhin Eiseigaku Zasshi* 10(6): 371-375.

- \* (169) Katz, J., and J. Kornblatt. 1962. Propionate metabolism by slices of mammary gland and liver of lactating rat. *J. Biol. Chem.* 237: 2466-2473.
- (170) Kaufmann, W., and K. Rohn. 1966. Gas chromatographic determination of volatile fatty acids in the rumen during various diets. *Z. Tierphysiol. Tierernachr. Futtermittelk.* 22(1): 1-8.
- (171) Kawada, N., and N. Hosoya. 1959. Influence of estradiol and toxohormone on the oxidation of fatty acids. *Seikagaku* 31: 734-738.
- (172) Kawakami, T., and K. Horiuchi. 1968. Seasonal changes of components and physical properties of apples. *Tokyo Nogyo Daigaku Nogaku Shuho* 1968: 12(3-4): 127-137.
- (173) Kern, J., et al. 1954. Organic acids formed during the heat sterilization of milk. *J. Dairy Sci.* 37: 904.
- (174) Kiely, M., and P. O'Driscoll. 1971. Lactic and volatile acids in beet sugar manufacture. I. Lactic and volatile acids in molasses. *International Sugar Journal* 73(869): 135-139.
- (175) Kimura, S., and K. Shioda. 1966. Dehydrated foods by freeze-drying. VII. Selection of antioxidants by indirect addition of antioxidant. 1. *Nippon Shokuhin Kogyo Gakkaishi* 13(3): 95-100.
- (176) Kirchner, J.G., and J.M. Miller. 1957. Volatile water-soluble and oil constituents of Valencia orange juice. *Jour. Agric. and Food Chem.* 5(4): 283-291.
- (176a) Kirk - Othmer Encyclopedia of Chemical Technology. 1967. 2nd. Ed. Volume 16.. Interscience Publishers, New York. pp. 554, 556.
- (177) Kitto, G.B., et al. 1962. Vitamin B<sub>12</sub>-coenzyme. *Sci. Cult.* 28: 295-302.
- (178) Klenk, E., and W. Bongard. 1952. Method for the chromatographic separation and determination of fragments formed on oxidative ozonide degradation of unsaturated fatty acids. *Hoppe-Seyler's Z. physiol. Chem.* 290: 181-198.

- (179) Knoop, F., and H. Jost. 1924. Einfluss einiger aliphatischer Substanzen auf Milchsäure- und Zuckerspiegel des Blutes. Ztschr. f. physiol. Ch. 141: 55-61.
- (180) Kooistra, J.A. and J. A. Troller. 1968. Food preservative compositions and method for inhibiting microbial growth in food. United States Patent 3 404 987.
- (181) Kosaki, T., et al. 1956. The influence of the administration of amino acids upon the distribution of porphyrin bodies in the tissue. IV. The influence of the administration of aminobutyric acid and related substances. Mie Med. J. 6: 47-53.
- (182) Kostal, J. 1940. Propionic acid in view of its physiological action. Chem. Zentr. 1941: I: 3250.
- (183) Kostanyan, G. G., et al. 1969. Polarographic determination of acrylic acid in commercial propionic acid. Arm. Khim. Zh. 1969: 22(11): 1044.
- (184) Kostanyan, G. G., et al. 1970. Chromatographic determination of the composition of a mixture of products obtained during butyric acid synthesis. Arm. Khim. Zh. 23(2): 134-139.
- (185) Kretovich, V. L., et al. 1951. Chromatographic determination of volatile fatty acids. Doklady Akad. S. S. S. R. 80: 409-412.
- (186) Kristofferson, T. 1963. Cheese flavor in perspective. Nanuf. Milk Prod. J. 54(5): 12-14.
- (187) Krylova, N. N., and K. Y. Bazarova. 1970. A study of the volatile fatty acid content of cooked sausages. Trudy Vsesoyuznyi Nauchno-Issledovatel'skii Institut Myasnoi Promyshlennosti 23: 26-31.
- (188) Kuhn, W., et al. 1958. Continuous gas chromatography. I. Procedure for continuous separation of a mixture of several components in two-phase countercurrent operation with temperature gradient. Helv. Chim. Acta. 41: 2135-2148.
- (189) Kuhn, W., et al. 1960. Continuous separation of multicomponent mixtures by gas chromatography. World Petrol. Congr., Proc. 5th N. Y. 1959: 45-53.
- (190) Kurtz, F. E., et al. 1959. Interrelations between pH, populations of Propionibacterium shermanii, levels of free fatty acids, and the flavor ratings of Swiss Cheeses. J. Dairy Sci. 42: 1008-1019.
- (191) Kuvaev, B. E., and N. S. Imyanitov. 1967. Determination of monobasic and dibasic acids by paper chromatography. Zavod. Lab. 33(11): 1382-1383.

- (192) Kuzdzal-Savoie, S., et al. 1971. Free fatty acids in Munster cheese.  
• Revue Laitieue Francaise. 290: 615.617.621.623.625.627 and 629.
- (193) Landes, D.R., and L.E. Dawson. 1971. Organic acid patterns  
in fresh and incubated whole egg products. Poultry Science  
50(2): 641-642.
- (194) Lang, K. and K.H. Bassler. 1953. Inhibition of acetoacetic acid  
formation by propionic acid. Klin. Wchnschr. 31: 675.
- (195) Langler, J.E., and E.A. Day. 1966. Quantitative analysis of the major  
free fatty acids in Swiss cheese. J. Dairy Sci. 49(1): 91-93.
- \* \*(196) Larson, P.S., et al.. 1956. Effect of chemical configuration on the  
edema-producing potency of acids, aldehydes, ketones, and alcohols.  
J. Pharmacol. Exptl. Therap. 116: 119-122.
- (197) Launer, H.F., and Y. Tomimatsu. 1953. Rapid accurate determination  
of carbohydrates and other substances with the dichromate heat-of-  
dilution method. Anal. Chem. 25: 1767-1769.
- (198) Lauryssens, M., et al. 1957. Metabolism of propionate 1-C<sup>14</sup> in the  
isolated udder of the cow. Arch. Internatl. Pharmacodyn. et Ther.  
109(1/2); 203-209.
- (199) Lebacq, J.P., et al. 1969. Thin-layer chromatographic separation of  
lower fatty acids (C<sub>2-9</sub>) in the form of p-bromoanilides and p-toluidid.  
Bull. Rech. Agron. Gembloux 4(1): 130-138.
- (200) Ledford, R.A. 1969. Direct gas-chromatographic measurement of  
acetic, propionic, and butyric acids in milk serum and aqueous ex-  
tracts of cheese. J. Dairy Sci. 52(7): 949-952.
- (201) Lefranc, L. 1922. Herstellung von Buttersaure und anderen aliphatischen  
Carbonsauren durch Vergaren von Zuckerlosungen. Canada Patent  
234 159 vom 19/8.
- (202) Lehmann, W., and K. Sahli. 1957. Determination of the lower fatty  
acids by Harper's chromatographic columns method. Proc. Intern.  
Dairy Congr. 14th, Rome 1956, 2: 314-322.
- (203) Lemeshek, K. 1969. Chemical substances for foodstuffs conservation.  
Moscow, Pishchevaya promyshlennost, 103 pp.
- (204) Leng, R.A., and E.F. Annison. 1963. Metabolism of acetate, propio-  
nate, and butyrate by sheep liver slices. Biochem. J. 86: 319-327.
- (205) Lenhart, J., and K.W. Cosens. 1949. Molds and bacteria curbed in  
immersion-freezing. Food Indust. 21(4): 442-444.

- (206) Levine, R.M., and E.W. Pelikan. 1961. Influence of experimental procedures and dose on the intestinal absorption of an onium compound, benzomethamine. *J. Pharmacol. Exptl. Therap.* 131: 319-327.
- \* (207) Lifson, N. 1957. Metabolism of C<sup>13</sup>-carboxyl-labeled propionate by the isolated, perfused dog gastrocnemius. *Amer. Jour. Physiol.* 188(2): 227-232.
- (208) Limantov, G.A. 1970. Improving the technology of Kavkazskii grating cheese. *Molochnaya Promyshlennost'* 31(10): 12-14.
- (209) Lindqvist, B., et al.. 1953. Determination of the volatile acids in cheese by paper chromatography. *Intern. Dairy Congr., Proc.* 13th (Hague) 3: 1250-1253.
- (210) Lipis, B.V., et al. 1967. Gas-liquid chromatographic study of the composition of volatile compounds in products of grape processing. *Sadovod. Vinograd. Vinodel. Mold*, No. 2: 28-32.
- (211) Lo, T.Y., and C.H. Wu, 1944. The peptization of rice proteins by organic acids and organic salts. *Chinese J. Sci. Agr.* 1: 173-178.
- (212) Long, C., and R.A. Peters. 1939. Brenztraubensaureoxydation im Hirn. V. Aus dem  $\alpha$ -Ketobuttersaurestoffwechsel erhaltene Hinweise. *Biochemical J.* 33: 759-773.
- (213) Lorber, V., et al., 1950. Conversion of propionate to liver glycogen in the intact rat, studied with isotopic propionate. *J. Biol. Chem.* 183: 531-538.
- (214) Luck, H. 1960. Influence of preservatives on hemin enzymes. II. A correlation between the preservative effect of organic acids and their catalase inhibition. *Z. Lebensm.-Untersuch. u.-Forsch.* 111: 190-198.
- \* (215) Luick, J.R. 1961. Synthesis of milk fat in the bovine mammary gland. *J. Dairy Sci.* 44: 652-657.
- (216) Luis, P., et al. 1968. Ultramicro detection of formic, acetic, and some other volatile organic acids. *Mikrochim. Acta* 1968(1): 56-60.
- (217) Lukac, S., and B. Bartonicek. 1969. Gas chromatographic estimation of liquid products of the radiolysis of solid malonic and succinic acids. *Chromatographia* 5: 192-196.



- (218) Lurje, H.S. 1925. Untersuchungen über die motorische Funktion des Dickdarms. I. Mitt. Die Einwirkung natürlicher chemischer Reizmittel auf die Bewegung des Dickdarms. Pflugers Arch. d. Physiol. 207: 269-278.
- (219) Lurje, H.S. 1926. Untersuchungen über die motorische Funktion des Dickdarms. IV. Mitt. Untersuchungen am überlebenden Dickdarm. Pflugers Arch. d. Physiol. 212: 64-76.
- (220) Maekawa, K, et. al. 1967. Gas chromatography of natural compounds. Mem. Ehime Univ., Sect. VI 12(3): 131-147.
- (221) Mahadevan, V., and L. Zieve. 1969. Determination of volatile free fatty acids of human blood. J. Lipid Res. 10(3): 338-341.
- \* (222) Manns, J.G., et al. 1967. Probable role of propionate and butyrate in control of insulin secretion in sheep. Amer. J. Physiol. 212(4): 756-764.
- \* (223) Manns, J.G., and J.M. Boda. 1967. Insulin release by acetate, propionate, butyrate, and glucose in lambs and adult sheep. Amer. J. Physiol. 212(4): 747-755.
- (224) Masoro, E.J., and S.S. Panagos. 1956. Effects of short chain fatty acids on hepatic acetate metabolism in cold exposure and fasting. Proc. Soc. Exptl. Biol. and Med. 93(2): 277-280.
- (225) Masoro, E.J., et al. 1957. The regulation of hepatic acetate-1-C<sup>14</sup> metabolism by short-chain fatty acids. Arch. Biochem. and Biophys. 68(2): 270-283.
- ⊙ (226) Masoro, E.J., and E. Porter. 1961. Propionic acid as a precursor in the biosynthesis of animal fatty acids. Jour. Lipid Res. 2(2): 177-181.
- (227) Mattick, L.R., and W.B. Robinson. 1960. Changes in the volatile acids during the baking of sherry wine by the Tressler baking process. Am. J. Enol. Viticult. 11: 113-116.
- (228) Maurer, H., and K. Lang. 1956. Effect of propionic acid on the metabolism of alloxan-diabetic rabbits. Klin. Wochschr. 34: 862-864.

- (229) Mazumder, R., et al. 1962. Metabolism of propionic acid in animal tissues. IX. Methylmalonyl coenzyme A racemase. J. Biol. Chem. 237: 3065-3968.
- (230) Mazumder, R., et al. 1963. Metabolism of propionic acid in animal tissues. X. Methylmalonyl coenzyme A mutase holoenzyme. J. Biol. Chem. 238: 50-53.
- \* \* (231) McAtce, J.W., et al. 1968. Utilization of rumen metabolites as energy sources in rats. Life Sci. 1968: 7(14)(Pt. 2): 769-775.
- (232) McRoberts, L.H. 1950. Report on mold inhibitors, propionates. J. Assoc. Offic. Agr. Chemists 33: 677-684.
- (233) McRoberts, L.H. 1951. Report on acetic and propionic acids in bread, mold inhibitors, propionates. J. Assoc. Offic. Agr. Chemists' 34: 284-296.
- (234) Meerov, I., et al. 1968. Lipids in the shells of high oil-yielding sunflower seeds. Maslozhir. Prom. 1968: 34(5): 12-14.
- \* \* (234a) The Merck Index. 1968. 8th Ed. Merck and Co. Rahway, New Jersey pp. 195, 873-4, 695, 1042.
- (235) Meyer, J., et al. 1952. The antituberculosis bacterial factors of cheese. Lait 32: 512-514.
- (236) Milch, R.A. 1961. Biochemical studies of osteoarthritis interactions of certain intermediary metabolites with collagen. Atti Congr. Lega Intern. Reumatismo, 10<sup>o</sup>, Rome 2: 409-412.
- \* (237) Mori, K. 1953. Production of gastric lesions in the rat by the diet containing fatty acid. Japanese Jour. Cancer Res. 44(4): 421-427.
- (238) Moriguchi, S., et al. 1961. Organic acids in soysauce. III. Composition of organic acids in wheat and wheat bran. Hakko Kogaku Zasshi 39: 293-297.
- (239) Morita, H., et al. 1969. Flavor components in mirin. II. Low volatility components. Hakko Kogaku Zasshi 47(5): 303-307.
- (240) Moriyama, S., et al. 1964. Putrefaction of foods. IV. Study on putrefaction products of fish stored in a refrigerator, a hot box, in a thermostat, and at room temperature. Shokuhin Eiseigaku Zasshi 5(5): 361-365.

- (241) Morr, C. V. et al., 1957. Some organic acids in raw and heated skim-milk. Jour. Dairy Sci. 40(8): 964-972.
- (242) Mossel, D. A. A., and A. S. deBruin. 1954. A gelatin liquefaction test for the screening of compounds used or proposed as inhibitors of bacterial proteolytic deterioration in foods. Jour. Microbiol. and Serol. 20(2): 233-240.
- (243) Mukerjee, H. 1959. Paper chromatography of volatile fatty acids. Anal. Chem. 31: 1284.
- (244) Naguib, M., and E. Beerbom. 1957. Chromatography of organic acids. Kiel. Milchwirtsch. Forschungsber. 9: 161-175.
- (245) Nakabayashi, T., et al. 1971. The foul smell of kaki-shibu(astringent extract of persimmon fruits). Journal of Food Science and Technology 18(1): 20-23.
- (246) Nakamura, M. 1968. Effect of fatty acids on Shigella. Proc. Mont. Acad. Sci. 1968: 28: 51-57.
- (247) Nakanishi, T., et al. 1965. Cheese flavors. VII. Flavor compounds in oryzae-processed cheese. Rakuno Kagaku no Kenkyu 14(6): A245-A254.
- (248) Nakanishi, T., et al. 1965. Flavor compounds in Gouda cheese. Rakuno Kagaku no Kenkyu 14(5): A213-A222.
- (248a) NAS/NRC. 1965. Chemicals used in Food Processing. Publication 1274. pp. 4, 10, 12, 183.
- (248b) NAS/NRC 1972. Comprehensive Gras Survey. Table 2, pp. 9, 27, 28, 38; Table 11, part A, pp. 3, 9 ; Table 13A, pp. 44, 133-135, 189.
- (249) Nemtseva, L. I., et al. 1966. Separate determination of low-molecular weight fatty acids in natural waters by gas-liquid chromatography. Hidrokhim. Mater. 41: 129-141.
- (250) Nichikova, P. R., et al. 1969. Analysis of aqueous solutions of C<sub>1-4</sub> low-molecular weight aliphatic acids by a gas-liquid chromatographic method. Neftepererab. Neftekhim. 1969(6): 45-47.
- (251) Nielsen, V. H. 1966. Making butter flavor better. Amer. Dairy Rev. 28(11): 50, 62-63.
- (252) Nikitin, G. A., et al. 1971. Chromatographic method of determining volatile acids using a densitometer. Mikrobiologiya 40: 575-578.

- (253) Obara, J., and H. Watase. 1957. Effects of administration of some carbohydrates and fatty acids upon the blood glucose level of goats. Bull. Natl. Inst. Animal Health 32: 373-377.
- (254) Obata, Y., and K. Matano. 1952. The flavor of katsuobushi. II. Identification of volatile-substances by paper chromatography. J. Agr. Chem. Soc. Japan 26: 184-185.
- (255) Obata, Y., and K. Matano. 1953. The odor of cuttlefish-liver oil. I. Identification of volatile acids and bases. J. Oil Chemists' Soc. 2: 112-114.
- (256) Oberdisse, K., and J.N. Paraskevopoulos. 1938. Über die spezifisch-dynamische Wirkung einiger stickstofffreier Substanzen an der isolierten Leber. Z. ges. exptl. Med. 102: 374-389.
- (257) Oerskov, E.R., et al. 1969. Influence of ruminal infusion of volatile fatty acids on milk yield and composition and on energy utilization by lactating cows. Brit. J. Nutr. 23(3): 443-453.
- (258) Oldfield, J.F.T. 1957. Formation of volatile organic acids in factory juices and the estimation of these acids by gas-liquid chromatography. Compt. rend. assemblee comm. intern. tech. sucriere, 10, London 1957: 56-66.
- \* (258a) OPD Chemical Buyers' Directory. 1970-1971. Schnell Publishing Co., Inc. New York. pp. 497, 1413.
- (259) Opienska-Blauth, J., et al. 1951. The partition paper chromatography of some organic acids. Ann. Univ. Mariae Curie-Sklodowka, Sect. D-Med 5(9): 221-275.
- \* (260) Oro, L., and A. Wretlind. 1961. Pharmacological effects of fatty acids, triolein, and cotton-seed oil. Acta Pharmacol. Toxicol. 18: 141-152.
- (261) Orskov, E.R., and D.M. Allen. 1966. Utilization of salts of volatile fatty acids by growing sheep. III. Effect of frequency of feeding on the utilization of acetate and propionate by young growing lambs. Brit. J. Nutr. 20(3): 509-517.
- (262) Osawa, S., et al. 1957. Mononucleotides of the cell nucleus. Jour. Gen. Physiol. 40(3): 491-513.
- (263) Osburn, O.L., et al. 1935. Die thermophile Gärung von Rubenbrei. J. Agric. prat. J. Agric. 50: 165-172.

- (277) Preonas, D.L. 1969. Modification of baked cream pie formulations to inhibit surface growth of *Staphylococcus aureus* MF31 without refrigeration. Dissertation Abstracts International. Section B. The Sciences and Engineering 30(3): 1189.
- (278) Prodanski, P. 1964. Some volatile aromatic components of cheese. *Prumysl Potravin* 15: 273-274.
- (279) Prodanski, P.G., and O.D. Dzhondzhorowa. 1969. Studies on some volatile substances and lipid fractions formed during ripening of the Bulgarian White pickled and Kachkaval cheese. *Milchwissenschaft* 24(12): 734-736.
- (280) Quastel, J.H., and A.H.M. Wheatley, 1933. Oxydation von Fettsauren in der Leber. *Biochemical J.* 27: 1753-1762.
- (281) Radtke, R., et al. 1966. Coffee aromas and their changes during storage of roasted coffee. III. Determination of low-volatile aroma compounds and volatile acids. *Z. Lebensm.-Untersuch. -Forsch.* 129(6): 349-359.
- (282) Raible, K. 1957. Effect of the initial concentration of yeast cells on the effectiveness of chemical preservatives in malt-wort. *Deut. Lebensm.-Rundschau* 53: 155-156.
- (283) Raible, K. 1957. Effect of age of yeast cultures on the effectiveness of chemical preservatives. *Deut. Lebensm.-Rundschau.* 53: 198-200.
- (284) Raible, K., and G. Busch. 1957. Studies on chemical conserving material I. The influence of the common preservatives on the multiplication of yeasts. *Zentr. Bakteriolog., Parasitenk., Abt. II* 110: 172-177.
- (285) Rao, S.V.S., and A.P. Valsan, 1963. Improvement in quality and storage life of pickled fish by means of propionic acid. *Indo-Pacific Fisheries Council, Proc.* 10(2): 315-317.
- (286) Raupp, G. 1959. Gas chromatographic separation of the lower fatty acids. *Angew. Chem.* 71: 284-285.
- (287) Raveux, R., and J. Bove. 1957. Separation and determination of carboxylic acids from C<sub>1</sub> to C<sub>6</sub> by partition chromatography on a silica column. I. General study. *Bull. soc. chim. France* 1957: 369-376.
- (288) Reagan, J.G., et al. 1971. Improved methods for determination of certain organic acids in pasteurized and unpasteurized liquid and frozen whole egg. *Journal of Food Science* 36(2): 351-354.

- (289) Rei, M. 1970. Lactic acid bacteria cultures in the production of sausage. *Myas. Ind. SSSR* 41(1): 37-39.
- (290) Reid, J. D. 1932. Die Desinfektionswirkung verschiedener organischer Sauren. *Amer. Jour. Hygiene* 16: 540-556.
- (291) Reid, R. L., and M. Lederer, 1951. Separation and estimation of saturated C<sub>2</sub>-C<sub>7</sub> fatty acids by paper-partition chromatography. *Biochem. J.* 50: 60-67.
- (292) Reshef, L., et al. 1967. Effect of propionate on lipogenesis in adipose tissue. *J. Lipid Res.* 8(6): 682-687.
- (293) Reshef, L., et al. 1967. Effect of propionate on pyruvate metabolism in adipose tissue. *J. Lipid Res.* 8(6): 688-691.
- (294) Rice, A. C., and C. S. Pederson, 1954. Factors influencing growth of *Bacillus coagulans* in canned tomato juice. I. Size of inoculum and oxygen concentration. *Food Research* 19: 115-133.
- (295) Richardson, L. R., and J. V. Halick, 1957. Feed spoilage. Heat-inhibiting activity of various compounds and commercial products. *Texas Agr. Exptl. Sta., Bull. No. 879*: 6 pp.
- (296) Ritter, W., and H. Haenni, 1960. Gas chromatography and its application in the dairy industry. II. Detection and determination of volatile fatty acids in dairy products and cultures. *Milchwissenschaft* 15(6): 296-302.
- (297) Rodel, W., et al. 1969. Meat flavour and natural ageing. II. The effect of meat constituents on flavour. *Nahrung* 13(6): 495-503.
- (298) Rogers, T. A., and M. Kleiber, 1956. The biosynthesis of milk-fat in the intact cow. *Biochim. et Biophys. Acta* 22(2): 284-292.
- (299) Rook, J. A. F., and C. C. Balch, 1961. The effects of intraruminal infusions of acetic, propionic and butyric acids on the yield and composition of the milk of the cow. *Brit. Jour. Nutrition* 15(3): 361-369.
- (300) Rook, J. A. F., et al. 1965. Further observations on the effects of intraruminal infusions of volatile fatty acids and of lactic acid on the yield and composition of the milk of the cow. *Brit. J. Nutr.* 19(1): 93-99.

- (301) Rosculet, G. 1969. Metabolic products of yeast fermentation: organic acids, isolation, separation, and characterization. Amer. Soc. Brew. Chem., Proc. 1969: 127-138.
- (302) Rubinstein, R., et al. 1969. In vivo dialysis of feces as a method of stool analysis. IV. Organic anion component. Clin. Sci. 37(2): 549-564.
- (303) Sandine, W.E., and P.R. Elliker. 1969. Flavour in fermented dairy products. Abstracts of Papers. American Chemical Society 158: MICR 29.
- (304) Sartory, R., et al. 1951. Propionic fermentation and acid-alcohol-resistant bacteria in hard cheeses. Bull. Acad. nat. med. 135: 708-710.
- (305) Sasaki, S., et al. 1969. Chemical studies on components of dried bonito, Katsuobushi. II. Volatile, neutral, noncarbonyl oxygenated compounds. Agr. Biol. Chem. 33(7): 1037-1041.
- (306) Schelhorn, M.v. 1953. Bacteriostatic and bactericidal action of preservatives. Arch. Mikrobiol. 19: 30-44.
- (307) Schluter, Th. 1932. Backen von Brot. Netherlands Patent 53 728 vom 16/11.
- (308) Schormuller, J., et al. 1960. The composition of the "tear fluid" of Emmentaler cheese. Z. Lebensm.-Untersuch. u. -Forsch. 112: 364-375.
- (309) Schormuller, J., and H. Langner. 1960. Quantitative determination of organic acids in foods. Z. Lebensm.-Untersuch. u. -Forsch. 113: 104-112.
- (310) Schulte, K.E., and J. Kirschner. 1951. Reaction kinetics of branched-chain fat acids. II. Identification of branched-chain fat acids by means of the esterification reaction. Fette u. Seifen 53: 456-457.
- (311) Semichon, L., and M. Flanzky. 1932. Untersuchung über die organischen Säuren der Weine, Traubenmoste und natürlichen oder vergorenen Fruchtsäfte. II. Die fluchtigen Fettsäuren. Ann. Science agronom. Francaise 2: 504-528.

- (312) Senel, S.H., and F.G. Owen. 1966. Relation of dietary acetate and lactates to dry matter intake and volatile fatty acid metabolism. *J. Dairy Sci.* 49(9): 1075-1079.
- (313) Shaburov, M.A., et al. 1970. Chromatographic determination of low-molecular-weight monocarboxylic acids and their butyl esters. *Neftepererab. Neftekhim.* 1970(4): 46-48.
- \* (314) Shcheptova, G.A. 1970. Hygienic standardization of the content of propionic acid and its sodium salt in water bodies. *Gig. Sanit.* 35(3): 96-98.
- (315) Shelley, R.N., et al. 1963. Determination of formic, acetic, propionic, and butyric acids by gas chromatography. *J. Assoc. Offic. Agr. Chemists* 46(3): 486-493.
- (316) Sherman, J.M., and R.H. Shaw. 1923. Die Propionsauregarung der Lactose. *Journ. Biol. Chem.* 56: 695-700.
- (317) Shibasaki, K., and S. Kimura. 1969. Food chemical studies on soybean phospholipids. III. Volatile fatty acids and alcohols from autoxidized phospholipids. *Journal of Food Science and Technology* 16(6): 277-281.
- (318) Shibata, F., and M. Umezu. 1961. Administration of acetic, propionic, and butyric acid alone and metabolic disturbances in ruminants. Proceedings of the 51st meeting of the Japanese Society of Veterinary Science, Tokyo, Japan, *Japanese Jour. Vet. Sci.* 23(Suppl.): 459-560.
- (319) Shiring, E.R., and A.T. Zagorul'ko. 1969. Products of the decomposition of sucrose during vaporization, and associated substances in ammonia water. *Sakh. Prom.* 43(4): 16-20.
- (320) Siigur, J., et al. 1969. Infrared spectra of acetic and propionic acids in the  $D_2O$ - $D_2SO_4$  system. *Reakts. Sposobnost Org. Soedin.* 6(2): 486-491.
- (321) Sinclair, W.B., and V.A. Jolliffe. 1960. Analytical methods for soluble carbohydrates and pectic substances in citrus fruits. *Conserve e deriv. agrumari* 9: 193-197.



- (322) Singer, I., et al. 1969. Effect of propionate and other organic anions on sodium transport across toad bladder. *Biochim. Biophys. Acta* 193(2): 430-443.
- (323) Singh-Verma, S.B. 1970. Application of propionic acid for preserving mixed feeding stuffs, industrially prepared, as well as cereals and maize. Results of microbiologic studies. *Zentralbl Bakteriologie* 125: 100-111.
- (324) Sjostrom, G., and S. Willart. 1954. Breakdown of fat in Swedish hard rennet cheese. *Svenska Mejeritidn.* 46: 277-280, 282-284, 287, 293-295, 297-298.
- (325) Smith, R.M., et al., 1967. Metabolism of propionate by sheep liver. Pathway of propionate metabolism in aged homogenate and mitochondria. *Biochem. J.* 104: 441-449.
- (326) Smith, R.M., et al. 1967. Metabolism of propionate by sheep-liver mitochondria. Evidence for rate control by a specific succinate oxidase. *Biochem. J.*, 104: 460-472.
- (327) Smouse, T.H., and S.S. Chang. 1967. A systematic characterization of the reversion flavor of soybean oil. *J. Amer. Oil Chem. Soc.* 44(8): 509-514.
- (328) Solomon, N.A., and S.Z. Lewin. 1955. Drop density, weight, volume, and time variations during chromatographic-column development. *Ann. N. Y. Acad. Sci.* 62: 93-104.
- (329) Somers, M., 1969. Volatile fatty acid clearance studies in relation to vitamin B<sub>12</sub> deficiency. *Aust. J. Exptl. Biol. Med. Sci.* 47(2): 219-225.
- (330) Stasch, A.R. 1958. Selection of liquid stationary phases for the separation of volatile mixtures by gas chromatography. *Bull. Georgia Acad. Sci.* 16: No. 3: 38-45.
- (331) Steinhauer, J.E., and L.E. Dawson. 1969. Quantitative determination of formic, acetic, propionic and butyric-acids in frozen whole eggs by gas-liquid chromatography. *J. Food Sci.* 34(4): 359-364.

- (332) Storgards, T., and B. Lindqvist. 1952. Different methods for demonstrating faulty fermentation in cheese. Svenska Mejeritidn. 44: 103-107.
- (333) Storry, J.E., and J.A.F. Rook. 1962. Effect of large intraruminal additions of volatile fatty acids on the secretion of milk constituents. Intern. Dairy Congr., Proc. 16th, Copenhagen, 1962 1: 64-70.
- \* (334) Storry, J.E., and J.A. Rook. 1965. Effect in the cow of intraruminal infusions of volatile fatty acids and of lactic acid on the secretion of the component fatty acids of the milk fat and on the composition of blood. Biochem. J. 96(1): 210-217.
- (335) Suomalainen, H., and E. Archimo. 1946. Determination of fatty acids of low molecular weight by the half-distillation method. Mitt. Lebensm. Hyg. 37: 173-178.
- (336) Suryanarayana Rao, S.V., and A.P. Valsan. 1962. Pickling of mackerels in tropical countries using propionic acid as preservative. Proc. Intern. Congr. Food Sci. Technol., 1st, London, 1962, 2: 221-226.
- (337) Suzue, M. 1926. Untersuchungen über die physikalisch-chemischen Eigenschaften der Zellmembran. III. Mitt. Über die Faktoren, die die Sedimentationsgeschwindigkeit der roten Blutkörperchen bestimmen. Journ. Biophysics 1: 259-269.
- (338) Syunitito Yosii. 1937. Beiträge zur Kenntnis der Fettsäureoxydation im Tierkörper. J. Biochemistry 26: 397-424.
- (339) Szonntag, E. 1959. Determination of the total and some more important volatile acids in cheese. Intern. Dairy Congr., Proc. 15th Congr. London 3: 1467-1473.
- (340) Takei, S., et al. 1934. Über die Riechstoffe des grünen Tees. III. Die sauren Stoffe im rohen Teeöl. Sci. Pap. Inst. physic. chem. Res. 25: Nr. 536/38.
- (341) Takeshita, R., et al. 1970. Qualitative and quantitative analysis of propionic acid from [in] foods by gas chromatography. Shokuhin Eiseigaku Zasshi 11(3): 143-146.
- (342) Tanaka, K., et al. 1967. The effects of organic and inorganic acids on the physical properties of dough. Hakkō Kagaku Zasshi 45(6): 566-569.

- (343) Tarantola, C. 1971. Composition of grappa in relation to the development of the volatile constituents of the pressed grape lees during storage and to the distillation technique. *Vini d'Italia* 13(73): 331-335.
- (344) Thompson, A.R. 1951. Volatile products of apples. I. Identification of acids and alcohols, *Australian J. Sci. Research B4*: 283-290.
- (345) Tizzano, A. 1946. The relation between bactericidal action and concentration of disinfectants. *Atti acad. Lincei, Classe sci. fis., mat. e nat.* [8]: 1: 225-231.
- \* (346) Tjan, G.H., and J.T.A. Jansen. 1971. Identification of acetic, propionic, and sorbic acid in bakery products by thin layer chromatography. *Journal of the Association of Official Analytical Chemists* 54(5): 1150-1151.
- \* \* (347) Tolnai, S., and J.F. Morgan. 1961. In vitro antitumor activity of fatty acids. III. Saturated monocarboxylic acids. *Can. J. Biochem. and Physiol.* 39: 713-719.
- (348) Tranger, M. 1963. A method for quantitative determination of lower fatty acids in serum by paper chromatography. *Vet. Arhiv* 33: 248-252.
- (349) Treadway, R.H. 1967. Manufacture of potato starch. *Starch: Chem. Technol.* 1967: 2: 87-101.
- (350) Trčka, V. 1960. Spermicidal activity of some substances. *Časopis Lékařů Českých* 89L 631-635.
- (351) Tripod, J., et al. 1955. Intermediary metabolites and the isolated rabbit heart. *Arch. intern. pharmacodynamie* 104: 121-136.
- (352) Turowska, M. 1960. Biacridine derivatives as indicators in titration of weak acids. *Chem. Anal.* 5: 815-821.
- (353) Ueda, R., et al. 1959. Organic acids in soysauce. I. Determination of the organic acids in soysauce. *Hakko Kogaku Zasshi* 37: 94-99.
- (354) Ueno, T., and T. Kuramochi. 1961. Semi-chemical soy sauce. VII. Volatile components produced by HCl hydrolysis of defatted soybean. *Nippon Nagei Kagaku Kaishi* 35: 454-458.

- (355) Ueno, T. 1961. Semi-chemical soy sauce. IX. Inhibitory effects of an acid hydrolyzate of defatted soybeans on yeast fermentation. *Nippon Nogei Kagaku Kaishi* 35: 1184-1189.
- (356) Ulyatt, M. J. 1965. The effects of intraruminal infusions of volatile fatty acids on food intake of sheep. *New Zealand J. Agr. Res.* 8(2): 397-408.
- (357) Umeh, E. O. 1970. Separation and determination of low-molecular weight straight-chain ( $C_1$ - $C_8$ ) carboxylic acids by gas chromatography of their anilide derivatives. *J. Chromatogr.* 51(2): 147-154.
- (358) Valsan, A. P., et al. 1961. Propionic acid as a preservative for cured fish products. *J. Sci. Ind. Research.* 20D: 351-354.
- (359) Vodovar, N., and J. Flanzy. 1967. Route of penetration of ingested lipids into the stroma of the villi. *C.R. Acad. Sci., Paris, Ser. D* 264(6): 850-853.
- (360) Vorbeck, M. L., et al. 1961. Volatile flavor of sauerkraut. Gas chromatographic identification of a volatile acidic off-odor. *J. Food Sci.* 26(6): 569-572.
- (361) Wagner, H. G., and F. A. Isherwood. 1961. Silica gel chromatography of organic acids from plant tissue. *Analyst* 86: 260-266.
- (362) Waligóra, B., and Z. Było. 1958. Potentiometric adsorption analysis as applied to determination of adsorption isotherms of chromatographic adsorbents. *Zeszyty Nauk. Uniw. Jagiel., Ser. Nauk Mat.-Przyrod., Mat., Fiz., Chem.* No. 4: 93-126.
- (363) Walker, G. H., et al. 1964. The detection and determination of propionic acid and sorbic acid in bread and flour confectionery. *J. Assoc. Public Analysts* 2(1): 2-4.
- (364) Walker, G. H., and M. S. Green. 1965. The determination of propionic acid in foods. *J. Assoc. Public Analysts* 3: 87-92.
- (365) Wang, K. T. 1960. Polyamide-impregnated paper chromatography. II. Aliphatic acids and amides. *J. Chinese Chem. Soc.* 7: 64-68.
- (366) Watanabe, K., and Y. Sato. 1969. Changes of meat fats by various processings. V. Volatile acidic compounds in heat-degraded pork fat. *Agr. Biol. Chem.* 33(10): 1411-1418.

- (367) Watson, J.R., and P. Crescuolo. 1970. Gas chromatographic separation and detection of C<sub>1</sub> to C<sub>3</sub> monocarboxylic acids as the p-substituted benzyl esters. *Journal of Chromatography* 52(1): 63-67.
- \*(368) Webb, J.L. 1950. The actions of metabolic substrates and inhibitors on the rabbit auricle. *Brit. J. Pharmacol.* 5: 87-117.
- (369) Wegener, W.S., et al. 1967. Propionate oxidation in *Escherichia coli*. *Arch. Biochem. Biophys.* 121(2): 440-442.
- (370) Wegener, W.S., et al. 1968. Propionate metabolism. II. Factors regulating adaptation of *Escherichia coli* to propionate. *Arch. Biochem. Biophys.* 123(1): 55-56.
- (371) Weston, R.H. 1966. Factors limiting the intake of feed by sheep. I. The significance of palatability, the capacity of the alimentary canal to handle digesta, and the supply of glucogenic substrate. *Australian J. Agr. Res.* 17(6): 939-954.
- \* \* (372) White, R.P., and F. E. Samson, Jr. 1956. Effects of fatty-acid anions on the electroencephalogram of unanesthetized rabbits. *Am. J. Physiol.* 186: 271-274.
- (373) Wierzchowski, J., and H. Witkowski. 1966. Paper-chromatographic determination of short-chain volatile fatty acids of fish. *Mitt. Geb. Lebensmittelunters. Hygs.* 57(2): 101-127.
- (374) Wierzbička, H., et al. 1967. Determination of acetate and propionate using propionate (acetate) kinase from *Propionibacterium shermanii*. *Acta Biochem. Pol.* 14(1): 1-6.
- (375) Willart, S., and G. Sjöström, 1957. Chromatographic determination of fatty acids as a means for examination of fermented cheese. *Svenska Mejeritidn.* 49: 623-629.
- (376) Willhalm, B., et al. 1966. Flavors. XII. The flavor of fresh strawberries. Identification of the volatile acids and other compounds. *Helv. Chim. Acta* 49(1): 65-67.
- (377) Wiseblatt, L. 1967. Reduction of the microbial populations in flours incorporated into refrigerated foods. *Cereal Chem.* 44(3): 269-280.
- (378) Wood, H.G., et al. 1957. Lactose synthesis. II. The distribution of C<sup>14</sup> in lactose of milk from the perfused isolated cow udder. *Jour. Biol. Chem.* 226(2): 1023-1034.

- (379) Yamada, K., and M. Ino. 1969. Studies on shoyu flavour by gas chromatography. II. Flavour components produced by shoyu koji microflora. *Seasoning Science* 16(5): 15-20.
- (380) Yamanishi, T., et al. 1966. Flavor of black tea. IV. Changes in flavor constituents during the manufacture of black tea. *Agr. Biol. Chem.* 30(8): 784-792.
- (381) Yamazaki, M. 1966. Volatile components of mulberry leaves by gas chromatography. II. Acidic and the phenolic components. *Nippon Nogei Kagaku Kaishi* 40(12): 456-460.
- (382) Yokotsuka, T. 1954. Flavoring substances in soy sauce. XIII. The fungistatic substances in brewed soy sauce and the increase of the fungistatic nature by heating the soy sauce. *J. Agr. Chem. Soc. Japan* 28: 114-118.
- (383) Young, J.W. 1964. The metabolism of acetate, propionate, and butyrate in young milk-fed calves. *Dissertation Abstr.* 25(4): 2135.
- (384) Young, J.W., et al. 1965. Metabolism of acetate propionate and butyrate in young milk-fed calves. *J. Dairy Sci.* 48(8): 1079-1087.
- (385) Yuch, M.H., and F.M. Strong. 1960. Some volatile constituents of cooked beef. *J. Agr. Food Chem.* 8: 491-494.
- (386) Zakowska, Z. 1970. Estimation of volatile acids in beet molasses. *Roczniki Technologii i Chemii Zywnosci* 19: 37-45.
- (387) Zbinovsky, V., and R.H. Burris. 1954. New techniques for adding organic acids to silicic acid columns. *Anal. Chem.* 26: 208-210.
- (388) Ziffer, J., and A.S. Gaffney. 1969. Inhibition of microbial growth in aldonic acid and aldionate compositions. U.S. Patent 3 483 003.

## SODIUM PROPIONATE

### Bibliography

- (389) Anand, J. C., et al. 1959. Prevention of fungal spoilage in mango pickle with chemical preservatives. *Indian J. Agr. Sci.* 29: No. 2-3: 181-184.
- (390) Anon. 1961. Sodium Propionate; exemption from requirement for tolerance. *Federal Register* 26: 2593-2594.
- (391) Anon. 1963. Cold-pack cheese foods; standards of identity. *Federal Register* 28: 4508.
- (392) Anon. 1969. Comparative investigations on the effect of sodium propionate and sodium acetate on abomasal secretion in sheep. *Izv. Inst. Srav. Patol. Zhivotn., Bulg. Akad. Nauk.* 1969: 13: 301-307.
- \* (393) Baile, C. A., and J. Mayer. 1969. Depression of feed intake of goats by metabolites injected during meals. *Amer. J. Physiol.* 217(6): 1830-1836.
- (394) Ban, T., and M. Hojo. 1969. Central action of gamma-butyrolactone and gamma-hydroxybutyrate (GHB). II. Effect of GHB and related agents on the cortical dendritic responses. *Jap. J. Pharmacol.* 19(1): 89-101.
- (395) Beckmann, F. 1934. Über die Permeabilität des Froschmuskels unter dem Einfluss von verschiedenen Substanzen des Muskelstoffwechsels. *Plügers Arch. ges. Physiol. Menschen Tiere* 234: 469-475.
- (396) Blondin, G. A., et al. 1969. Mechanism of mitochondrial swelling. II. Pseudoenergized swelling in the presence of alkali metal salts. *Arch. Biochem. Biophys.* 129(1): 158-172.
- (397) Brachfeld, B. A. 1969. Antimicrobial food additives. *Bakers' Digest* 43(5): 60-62 & 65.
- (398) Childers, E., and G. W. Struthers. 1955. Infrared evaluation of sodium salts of organic acids. *Anal. Chem.* 27: 737-741.
- (399) Cornwell, R. T. K. 1951. Protective coating composition for hams. *U. S. Patent* 2 558 042.
- (400) Dalby, G., and H. C. Fisher. 1965. Emulsifier for cake batter. *U. S. Patent* 3 164 476.
- (401) Daugerts, R. 1965. Ketosis in cows and its cure with sodium propionate. *Tr. Lab. Biokhim. Fiziol. Zhivotn. Inst. Biol., Akad. Nauk Latv. SSR* 4: 163-166.

- (402) Dorello, U., et al. 1952. The in vitro action of some antibiotics and antiseptics on pathogenic organisms. *Boll. oculist.* 31: 281-290.
- (403) Dorello, U., et al. 1952. In vitro activity of sodium propionate toward various more common bacteria of ocular pathology. *Ann. ottal. e clin. ocul.* 78: 137-146.
- (404) Dowden, B. F., and H. J. Bennett. 1965. Toxicity of selected chemicals to certain animals. *J. Water Pollution Control Federation* 37(9): 1308-1316.
- (405) Eeckhaut, R. G. 1970. A new antiseptic in pork products manufacture. *Fermentatio* 66(1): 34-38.
- (406) Flatt, W. P., et al. 1958. Influence of purified materials on the development of the ruminant stomach. *Jour. Dairy Sci.* 41(1): 1593-1600.
- (407) Furchgott, R. F., and E. Shorr, 1946. Sources of energy for intestinal smooth muscle contraction. *Proc. Soc. Exptl. Biol. Med.* 61: 280-286.
- \* (407a) Furia, T. E. 1968. Handbook of Food Additives. The Chemical Rubber Co. Press. Cleveland, Ohio. pp. 159-163.
- (408) Gerlsma, S. Y. 1970. Gelatinizing temperatures of starch, as affected by polyhydric and monohydric alcohols, phenols, carboxylic acids, and some other additives. *Staerke* 22(1): 3-9.
- (409) Gompertz, D., et al. 1967. Metabolic precursors of methylmalonic acid in vitamin B<sub>12</sub> deficiency. *Clin. Chim. Acta* 18(2): 197-204.
- (410) Gonthier, B., and J. Mocotte. 1970. Biocidal preservatives for fish. *German* 1 921 670.
- (411) Graham, W. D., et al. 1954. Chronic toxicity of bread additives to rats. *J. Pharm. and Pharmacol.* 6: 534-545.
- \* \* (412) Graham, W. D., and H. C. Grice. 1955. Chronic toxicity of bread additives to rats. II. *J. Pharm. and Pharmacol.* 7: 126-134.
- \* (413) Hara, S., et al. 1963. Pharmacological and toxic actions of propionates. Examination of general pharmacological actions and toxicity of sodium and calcium propionates. *Tokyo Ika Daigaku Zasshi* 21(3, 4): 261-302.
- (414) Hendrickx, H., and A. de Vleeschauwer. 1952. Prevention of mold development on cheese rind. *Mededel. Landbouwhogeschool en Opzockingsstas. Staat Ghent* 17: 669-677.



- ✱ (415) Heseltine, W.W., and L.D. Galloway. 1951. Antibacterial properties of sodium propionate. *J. Pharm. Pharmacol.* 3: 581-585.
- (416) Heseltine, W.W. 1952. Sodium propionate and derivatives as bacteriostatics and fungistatics. *J. Pharm. Pharmacol.* 4: 577-581.
- (417) Hill, C.H. 1952. Propionic, inhibition of growth of *Streptococcus faecalis* by sodium propionate. *J. Biol. Chem.* 199: 329-332.
- (418) Huang, E., and J.G. Armstrong. 1970. The effect of microbial inhibitors on the storage life of Cottage cheese exposed to ambient temperatures. *Canadian Institute of Food Technology Journal* 3(4):157-161.
- (419) Hyson, A.M. 1959. Odor-controlled sodium propionate-sodium dipropionate compositions for use in baking. U.S. Patent 3 008 986.
- (420) Jackson, C.R. 1965. Laboratory evaluation of fungicides for control of some fungi found on peanuts. *Plant Disease Reprtr.* 49(11): 928-931.
- (421) Jalil, A., et al. 1963. Effect of metallic contamination and storage temperatures on keeping quality of khoa. *Agra Univ. J. Res.* 12: Pt. 1: 13-23.
- (422) Jendrassik, L., and H. Tangl. 1925. Die Atropinhemmung und die Wirkung einiger organischer Säuren am Darm. *Biochem. Ztschr.* 159: 337-350.
- ✱ (422a) Joint FAO/WHO Expert Committee on Food Additives. 1962. Evaluation of the toxicity of a number of antimicrobials and antioxidants. Sixth Report of the Joint FAO/WHO expert committee on food additives. Report No. 31. pp. 84-86.
- (423) Kashima, T. 1954. Titration of nonaqueous solutions of drugs and chemicals. III. The determination of sodium salts of carboxylic acids in glacial acetic acid. *Bull. Natl. Hyg. Lab.* No. 72: 145-150.
- (424) Keeney, E.L. 1943. Fungistatic and fungicidal effect of sodium propionate on common pathogens. *Bull. Johns Hopkins Hosp.* 73: 379-390.
- (425) Kerewala, D.N., and G.S. Siddappa. 1963. Fruit toffees. I. Effect of addition of glucose on texture and of processing temperature on retention of ascorbic acid and carotene in mango toffee. *Food Sci.* 12(8): 221-223.
- (426) Ketch, F.M., and R.L. Barton. 1970. Nonrefrigerated ready-to-bake pastry product. U.S. Patent 3 492 127.

- (427) Landsteiner, K., and J. van der Scheer. 1927. Einfluss organischer Salze auf die Wirkung der Tyrosinase. Proc. Soc. for exptl. Biology and Medicine 24: 692-693.
- (428) Liang, Y. T., et al. 1967. Absorption and utilization of volatile fatty acids by the young calf. J. Dairy Sci. 50(7): 1153-1157.
- (429) Luthman, J. 1968. The effect of propylene glycol on blood glucose and non-esterified fatty acids (NEFA) in ruminants. Nord. Vet. Med. 20(6): 338-343.
- (430) Macleod, G. K., and J. R. Luick. 1955. Acetate metabolism in ketosis of dairy cattle. Proc. 36th Ann. Meeting Western Div. Am. Dairy Sci. Assoc. 1955: 73-78.
- (431) Mariat, F., and M. Vieu. 1956. Fungistatic action of sodium propionate and of sodium dodecyl sulfate, and their synergism by combination. Ann. inst. Pasteur 91: 678-683.
- (432) McCullough, M. E., et al. 1969. Sodium acetate and sodium propionate as additives to all-in-one silage rations for milk production. J. Dairy Sci. 52(1): 1605-1608.
- (434) Medway, W., et al. 1957. The physiological effect of massive doses of mineral salts in pigs and chickens. Rept. N. Y. State Vet. Coll. Cornell Univ. 1955-1956: 22.
- \* (435) McManus, F. C. 1965. Stimulation of gastric acid secretion by acetate. Proc. Soc. Exptl. Biol. Med. 119(2): 473-476.
- (436) Molnar, S., et al. 1970. Fat metabolism studies in newborn calves with carbon-14-labeled substances. 1. Distribution of the carbon-14 activity in newborn calves 24 hours after oral administration of the carbon-14-labeled substances. Z. Tierphysiol., Tierernaehr. Futtermittelk 26(3): 121-132.
- (437) Netherland Appl. 6 512 060. 1966. Spread cheese.
- (438) Nicholson, J. W. G., et al. 1962. Addition of buffers to ruminant rations. II. Additional observations on weight gains, efficiency of gains, and consumption by steers of all-concentrate rations. Can. J. Animal Sci. 42: 75-81.
- (439) Nicholson, J. W. G., et al. 1962. The addition of buffers to ruminant rations. III. The effect of additions of sodium bicarbonate, sodium propionate, limestone and cod liver oil on apparent digestibility and nitrogen retention of an all-concentrate ration. Canadian Jour. Animal Sci. 42(1): 82-87.

- (440) Ochoa, A.G., and J. Ruiloba. 1944. Acción del propionato de sodio. Rev. Inst. Salubridad y Enferm. Trop. 5(2): 83-86.
- (441) Olivier, C., and M.T. Nguyen, 1959. Preservation of fish and animal meal. French Patent 1 253 279.
- \* (442) Pace, D.M., et al. 1967. Effects of sodium salts of certain monocarboxylic acids on established cell lines. Can. J. Biochem. 45(1): 81-88.
- (443) Park, H.S., et al. 1970. Survival of *Salmonella typhimurium* in cold-pack cheese food during refrigerated storage. Journal of Milk and Food Technology 433(9): 383-388.
- (444) Peeters, G., et al. 1953. The metabolism of propionic, butyric, and  $\beta$ -hydroxybutyric acids in the perfused isolated cow mammary gland. Arch. intern. pharmacodynamie 95: 153-171.
- (445) Pomerant, C.M., and C.D. Leake. 1954. Short-term cultures for drug assays. Ann. N.Y. Acad. Sci. 58: 1110-1128.
- (446) Raible, K. 1957. Effect of sub-threshold concentrations of chemical preservatives on the growth and fermentation of yeast in malt-wort. Deut. Lebensm.-Rundschau 53: 60-64.
- (447) Rakshy, S.E.S.E., and S. Abou-Donia. 1969. The inhibitory effect of Myprozine on moulds isolated from cheese-ripening rooms. Indian Journal of Dairy Science 22(2): 114-116.
- (448) Richards, C.R. 1958. Some observations on the use of sodium propionate for the prevention of ketosis in dairy cattle. Jour. Dairy Sci. 41(1): 1580-1585.
- (449) Richardson, L.R. and J.V. Halick. 1957. Feed spoilage. Heat-inhibiting activity of various compounds and commercial products. Texas Agr. Expt. Sta., Bull. No. 879: 6 pp.
- (450) Richou, R., and H. Richou. 1955. Antagonistic action of sodium propionate toward staphylococci. Compt. rend. soc. biol. 149: 943-945.
- \* (451) Romano, C. 1955. The biological activity of certain acids, ketones, and aliphatic esters. Arch. sci. biol. 39: 304-307.
- (452) Ruckebusch, Y., et al. 1967. Variations in blood constituents in relation to alimentary behavior in sheep. C.R. Soc. Biol. 161(2): 350-354.

- (453) Salunkhe, D.K., et al. 1959. Effects of  $\gamma$  radiation dose, rate, and temperature on the taste preference and storage life of certain fruits, vegetables, and their products. *Proc. Am. Soc. Hort. Sci.* 74: 414-422.
- (454) Sarra, C., et al. 1963. Synthesis of the fatty acids after intraruminal and intramammary administration of sodium propionate. *Ann. Fac. Med. Vet. Torino* 13: 119-125.
- (455) Savich, A.L., and C.F. Jansen. 1958. Color retention in meat. U.S. Patent. 2 830 907.
- (456) Schelhorn, M.v. 1951. Preservatives. V. Comparative effectiveness of preservatives. *Deut. Lebensm. Rundschau* 47: 16-18.
- (457) Schirmer, R., and W. Germans, 1954. Effect of sodium propionate in diseases of exterior eye. *Klin. Monatsbl. Augenh.* 125: 358-361.
- \* (458) Schmidt, G.H., and L.H. Schultz. 1958. Effect of feeding sodium propionate on milk and fat production, roughage consumption, blood sugar, and blood ketones of dairy cows. *J. Dairy Sci.* 41:169-175.
- (459) Schultz, L.H. 1954. Use of sodium propionate in the prevention of ketosis in dairy cattle. *Proc. Cornell Nutrition Conf. Feed. Manuf. Ithaca* 1954: 76-81.
- (460) Schwartzman, G. 1960. Separation and identification of sodium salts of  $C_2$  to  $C_5$  volatile fatty acids by paper chromatography. *J. Assoc. Offic. Agr. Chemists* 43: 428-430.
- \* \* (461) Serchi, G. 1954. Fungicidal, fungistatic, and antifermentative properties of some organic compounds. I. Sodium propionate. *Chimica* 9: 149-156.
- (462) Shchepetova, G.A. 1970. Hygienic standardization of the content of propionic acid and its sodium salt in water bodies. *Gig. Sanit.* 35(3): 96-98.
- (463) Smith, D.L.T. 1957. Poisoning by sodium salt - A cause of eosinophilic meningoencephalitis in swine. *Amer. Jour. Vet. Res.* 18(69): 825-850.
- (464) Spurgeon, K.R., et al. 1970. Develop low-fat dairy spread. *American Dairy Review* 32(9): 48 & 50.
- (465) Steger, H., et al. 1969. Effect of continuous intravenous and intraruminal infusions of volatile fatty acids on the feed consumption of dairy cows. *Arch. Tierernaehr.* 19(9): 641-649.

- (466) Steimetz, E. P. 1959. Application of the bioelectronics to chemical analysis. *Chim. anal.* 41: 321-331.
- (467) Stein, J. F., et al. 1958. Preliminary study on the use of sodium propionate with animal fat in dairy cattle rations. *Proc. Western Div. Am. Dairy Sci. Assoc.* 39: 5 pp.
- (468) Trenkle, A., and K. V. Kuhlemeier. 1966. Relation of rumen volatile acids, blood glucose, and plasma nonesterified fatty acids in sheep. *J. Anim. Sci.* 25(4): 1111-1115.
- (469) Umezu, M., et al. 1954. Ketosis in ruminants. 1. Production of experimental ketosis in goats by the administration of butyric acid. *Nippon Juigaku Zasshi* 16: 281-289.
- (470) Vaisey, E. B. 1954. A note on the control of dun. *Fisheries Research Board Can., Progr. Repts. Atlantic Coast Stas.* No. 58: 32.
- (471) Venkataraman, S., et al. 1967. Effect of propionate on the induction of vitamin B<sub>12</sub> deficiency in chicks and rats. *J. Nutr.* 93(2): 131-134.
- (472) Vicher, E. E., et al. 1968. The effect of sodium propionate and sodium caprylate on the fatty acid content of *Trichophyton rubrum*. *Mycopathol. Mycol. Appl.* 35(3/4): 208-214.
- (473) Williams, E. L., et al. 1969. Dietary depletion of vitamin B<sub>12</sub> and the excretion of methylmalonic acid in the rat. *Brit. J. Nutr.* 23(2): 343-352.
- (474) Young, J. A., and G. Schwartzman. 1963. Separation and identification of sodium salts of acetic, propionic, butyric, and valeric volatile fatty acids by paper chromatography. *J. Assoc. Offic. Agr. Chemists* 46: 483-485.
- \* (475) Young, J. A., et al. 1965. Separation and identification of sodium salts of acetic, propionic, butyric, and valeric acids by paper chromatography. *J. Assoc. Offic. Agr. Chemists* 48(3): 622-624.

## CALCIUM PROPIONATE

### Bibliography

- (476) Anon. 1940. Retarding mold in cheese. *Nation. Butter and Cheese Jour.* 31(8): 12.
- (477) Anon. 1963. Cold-pack cheese foods; standards of identity. *Federal Register* 28: 4508.
- (478) Anon. 1968. Current food additives legislation. - Netherlands. *Current Food Additives Legislation* 1968(116): 7.
- (479) Babel, F. J., and B. W. Hammer. 1939. Flavor contributors of swiss-type cheese. *Food Res.* 4(1): 81-85.
- (480) Balch, C. C., et al. 1967. The effect on milk yield and composition of adding the calcium salts of acetic, propionic, butyric and lactic acids to the diets of dairy cows. *J. Dairy Res.* 34(3): 199-206.
- (481) Barrett, F. 1970. Extending the keeping quality of bakery products. *Bakers' Digest* 44(4): 48-49 & 67.
- (482) Behrens, B., and J. Wajzer. 1935. Der lokale Reizeffekt von Calciumsalzen in Beziehung zu ihrer Dissoziation. *Biochem. Z.* 264: 120-125.
- (483) Belasco, J. 1959. Futtermittel für Wiederkäuer. *Austrian Patent* 2 965 488
- (484) Bergman, M. 1955. The determination of calcium propionate. *Chem. Tech.* 7: 483.
- (485) Blinc, M., et al. 1956. Propionic acid bacteria as inhibitors in the prevention of ropiness in bread. *Brot. u. Gebäck* 10: 66-68.
- (486) Bolin, H. R., et al. 1967. Effectiveness and stability of various antimicrobial agents on dates. Presented at the 27th Annual Meeting, Inst. of Food Technologists, May 14-19, 1967, Paper 123.
- (487) Bour, R. C. 1958. Wässerige Dispersionen für Bäckereizwecke. *Austrian Patent* 2 932 574
- (488) Brachfeld, B. A. 1969. Antimicrobial food additives. *Bakers' Digest* 43(5): 60-62 & 65.
- (489) Bray, R. W., and F. Hanning. 1949. Mold control in stored cured pork. *Jour. Animal Sci.* 8(4): 495-500.
- (490) Canuti, A. 1964. Analytische Untersuchung einiger kürzlich zugelassener Zusatzstoffe. *Boll. Lab. chim. provinc.* 15: 13-22.

- (491) Cole, D. J. A., et al. 1968. The effect on performance and bacterial flora of lactic acid, propionic acid, calcium propionate and calcium acrylate in the drinking water of weaned pigs. *Vet. Rec.* 83(18): 459-463.
- (492) Cook, R.M. 1970. Utilization of volatile fatty acids in ruminants. II. Comparison of the metabolism of acetate, propionate and butyrate when injected into the jugular vein, portal vein or the rumen. *Biochim. Biophys. Acta.* 201: 91-100.
- (493) Coolhaas, C. 1926. Thermophilic fermentation processes. *Ned. Tijdschr. Hyg. Microbiol. Serol.* 1: 338-350.
- (494) Corse, D.A. et al. 1970. Propionate utilization by pregnant, lactating, and spontaneously ketotic cows. *J. Dairy Sci.* 53: 740-746.
- (495) Dalby, G. 1949. pH control in bread and cake baking. *Bakers Digest* 23(1): 27-28, 38.
- (496) Diemair, W., and A. Schulz. 1968. Concerning knowledge of the prevention of mold-formation in sliced bread by sorboylpalmitate. *Ztschr. für Lebensmittel-Untersuchung und -Forschung* 138(1): 39-43.
- (497) Drews, E. 1955. Paper chromatography of bread and baked goods. II. Determination of volatile acids. *Brot u. Gebäck* 9: 45-47.
- (498) Drews, E. 1956. Zur Bestimmung von Calciumpropionat im freigeschobenen Roggenbrot. *Brot u. Gebäck* 10: 156.
- (499) French Patent 815 783, 815 784 u. 815 785, sämtlich vom 31/12 1936. 1936. Brotschimmelverhütung.
- (500) Greenwood, D. A., et al. 1945. Factors and processes influencing the keeping quality of bacon. *Proc. Inst. Food Tech.* 1945: 58-71.
- (501) Gross, H. 1968. Research with a pilot scale continuous breadmaking unit. VI. Buffering effects of calcium propionate in no-flour brews. *Cereal Science Today* 13(9): 346-348 & 358.
- (502) Hara, S., et al. 1963. Pharmacological and toxic actions of propionates. Examination of general pharmacological actions and toxicity of sodium and calcium propionates. *Tokyo Ika Daigaku Zasshi* 21(3, 4): 261-302.
- (503) Harshbarger, K.E. 1942. Report of a study on the toxicity of several food-preserving agents. *J. Dairy Sci.* 25: 168-174.
- (504) Hsia, Y.E., et al. 1969. Defective propionate carboxylation in ketotic hyperglycinaemia. *Lancet* 1: 757-758.

- (505) Iglauer, A. 1949. Konservieren von Gurken und Gemuse durch Einsauern (Einsalzen). Austrian Patent 169 092 vom 17/11 1949.
- (506) Kamaletdinov, A. Z. 1958. The preservation of bread to be used by mountain climbers during high-mountain climbing. Trudy L'vov Inst. Fiz. Kul'tury 3: 251-263.
- (507) Kamaletinov, A. Z. 1961. Effect of propionic acid and calcium propionate on the growth of molds. Mikrobiol. Zh., Akad. Nauk Ukr. RSR 23: No. 4: 19-22.
- (508) Kay, H. W. 1968. New international draft standards for processed cheese products. Milchwissenschaft 23 (1): 6-14.
- (509) Kiermeier, F. 1952. Food as ion exchange. I. Reaction of calcium acetate and calcium propionate in aqueous flour suspensions. Z. Lebensm.-Untersuch. u. -Forsch 94: 221-227.
- (510) Kiermeier, F. 1953. Lebensmittel als Ionenaustausch. 3. Mitt. Über die Reaktionen von Calciumpropionat in wasserigen Getreidestarkesuspensionen. Z. Lebensmittel-Unters. u. -Forsch. 97: 271-275.
- (511) Kimura, S., and K. Shioda. 1966. Dehydrated foods by freeze-drying. VII. Selection of antioxidants by indirect addition of antioxidants. I. Nippon Shokuhin Kogyo Gakkaishi 13(3): 95-100.
- (512) Kooistra, I. A. and J. A. Troller. 1968. Food preservative compositions and method for inhibiting microbial growth in food. U. S. Patent 3 404 987.
- (513) Kooistra, J. A. 1966. Low moisture content pastry dough compositions. U. S. Patent 3 294 547.
- (514) Lamprecht, F. 1955. Combating bread mold, especially Trichosporon variable. Brot u. Gebäck 9: 26-30.
- (515) Lebenbaum, M. 1936. Verhindern der Schimmelbildung in organischen Stoffen. Danish Patent 54 482 vom 22/12 1936.
- (516) Lloyd, D. 1968. Propionate assimilation in the flagellate Polytomella caeca. An inducible mitochondrial enzyme system. Biochem. J. 109: 897-907.
- \* (517) Lorber, V., et al. 1950. Conversion of propionate to liver glycogen in the intact rat, studied with isotopic propionate. Jour. Biol. Chem. 183(2): 531-538.



- (518) Macy, H., and J.C. Olson. 1939. Preliminary observations on the treatment of parchment paper with sodium or calcium propionate. J. Dairy Sci. 22: 527-534.
- (519) Macy, H. 1942. Mold inhibitors for food products. Assoc. Food Drug Officials U.S., Quart. Bull. 6: 9-12.
- (520) McRoberts, L.H. 1951. Report on acetic and propionic acids in bread, mold inhibitors, propionates. J. Assoc. Offic. Agr. Chemists 34: 284-296.
- (521) McRoberts, L.H. 1953. Report on acetic and propionic acid mold inhibitors in bread. J. Assoc. Offic. Agr. Chemists 36: 769-781.
- (522) Melnick, D., et al. 1956. Sorbinsäure als ein fungistatisches Agens für Nahrungsmittel. II. Mitt. Wirksamkeit von Sorbinsäure zum Schutz von Kuchen. Food Res. 21: 133-146.
- (523) Milatovic, L., and I. Bojanic. 1966. Influence of certain substances in reducing the thermophiles *Bacillus subtilis* and *B. messentericus* in bread. Presented at the 51st Annual Meeting, American Association of Cereal Chemists, April 12-15, 1966, Paper 71.
- (524) Milatovic, L., et al. 1966. Bread bacterial combinations. Kem. Ind. 15(12): 773-787.
- (525) Milner, M., et al. 1947. Grain-storage studies. VII. Influence of certain mold inhibitors on respiration of moist wheat. Cereal Chem. 24: 507-517.
- (526) Momtaz El-Gindy, M., and E. Drews. 1957. Paper-chromatographic identification of some preservatives in baked products. Brot u. Gebäck 10: 224-228.
- (527) Morrow, G., et al. 1969. Propionate metabolism in cells cultured from a patient with methylmalonic acidemia. Pediat. Res. 3: 217-219.
- \* (528) Nakao, K., et al. 1968. Utilization of propionate for heme synthesis in human bone marrow cells in vitro. J. Lab. Clin. Med. 72: 958-965.
- (529) Nury, F.S., et al. 1960. Praservierende Wirkung einiger antimikrobieller Agentien auf getrocknete Früchte von hohem Feuchtigkeitsgehalt. Food Technol. 14: 113-115.

- (530) O'Leary, D.K., and R.D. Kralovec. 1941. Development of *B. mesentericus* in bread and control with calcium acid phosphate or calcium propionate. *Cereal Chem.* 18: 730-741.
- (531) Olson, J.C., and H. Macy. 1942. Some observations on the treatment of parchment wrappers with sodium and calcium propionates for the inhibition of mold growth on the surface of butter. *Assoc. Southern Agr. Workers, Proc. Ann. Convention* 43: 122.
- (532) Olson, J.C. Jr., and H. Macy. 1946. Propionic acid, sodium propionate and calcium propionate as inhibitors of mold growth. II. Studies pertaining to the active agent responsible for the inhibitory effect of the propionates. *Jour. Dairy Sci.* 29(3): 173-180.
- (533) Orskov, E.R., et al. 1966. Utilization of salts of volatile fatty acids by growing sheep. II. Effect of stage of maturity and hormones implantation on the utilization of volatile fatty acids as sources of energy for growth and fattening. *Brit. J. Nutr.* 20(2): 307-315.
- (534) Orton, E.C. 1946. Dipping trials in drying fruit. *Australian Dried Fruits News* 21: 5.
- (535) Pette, J.W. 1953. Flavor development during cheese ripening. *Proc. 13th Intern. Dairy Congr.* 2: 557-564.
- (536) Pomeranz, Y., and L. Adler. 1959. Effectiveness of sorbic acid in inhibition of microbiological spoilage of bread. *Bull. Res. Council Israel Sect. C. Technol.* 6(4): 220-226.
- (537) Pozhogina, P.M., and V.V. Ruus. 1968. Method of preserving fat-containing products. *USSR Patent* 225 680.
- (538) Ramos, E.H., et al. 1969. Propionate metabolism in yeast. *Biochim Biophys Acta* 177: 671-673.
- (539) Reisener, H.J., et al. 1969. Propionate metabolism in the host-parasite complex of *Puccinia graminis*. *Naturwissenschaften* 56: 571.
- (540) Richardson, K.C. 1970. Chemical preservatives in foods in Australia. *Food Preservation Quarterly* 30(1): 2-8.
- (541) Ruczaj, Z., et al. 1969. Propionate and acetate kinase in *Streptomyces*. *Bull. Acad. Pol. Sci.* 17: 531-533.

- (542) Seiler, D.A.L. 1964. Application of fungicides in baking of bread and cakes. Int. Symp., Food Microbiol., 4th, Goteborg, Swed. 1964: 211-220.
- (543) Seto, K., et al. 1970. Metabolism of volatile fatty acids in rumen epithelium of ruminates. VII. Propionate metabolism and its antiketogenic action. J. Jap. Biochem. Soc. 42: 120-129.
- (544) Sieburg, E., and A. Kesler. 1923. Die Erhöhung der Calciumionen im menschlichen Serum nach intravenöser Zufuhr von Kalksalzen. Arch. f. exp. Pathol. u. Pharmak. 96: 180-190.
- (545) Slanina, L. 1970. Therapy of simple dysfunctions of the rumen in ruminants. Deut. Tieraerztl. Wochenschr. 77(1): 1-5.
- \* (546) Smyth, H.F. Jr., et al. 1969. Range-finding toxicity data. VII. Amer. Ind. Hyg. Ass. J. 30(5): 470-476.
- (547) Stern, J.S., et al. 1970. Are propionate and butyrate physiological regulators of plasma insulin in ruminants? Amer. J. Physiol. 219: 84-91.
- (548) Thompson, J.W., and E.R. Sherwin. 1966. Investigation of antioxidants for polyunsaturated edible oils. J. Amer. Oil Chemists' Soc. 43(12): 683-686.
- (549) Valette, G., and H. Huidobro, 1957. Die Wirkung organischer Calciumsalze auf die Sensibilität des isolierten Meerschweinchendarmes gegenüber Histamin. C R hebdom. Séances Acad. Sci. 245: 2093-2095.
- (550) Van Wieren, W. H. 1966. Making soft unripened curd cheeses. U. S. Patent 3 232 768.
- (551) Waitman, R. H., and J.W. Hoos. 1968. Ready-to-serve gelatin composition. U.S. Patent 3 367 784.
- (552) Wegener, W.S., et al. 1969. Propionate metabolism. V. The physiological significance of isocitrate lyase during growth of E. coli on propionate. Arch. Biochem. 129: 545-553.
- (553) Weidemann, M.J., et al. 1969. Acceleration of gluconeogenesis from propionate by DL-carnitine in the rat kidney cortex. Biochem. J. 111: 69-81.

- (554) Weidemann, M.J., et al. 1970. Gluconeogenesis from propionate in kidney and liver of the vitamin B<sub>12</sub>-deficient rat. Biochem. J. 117: 177-181.
- (555) Young, J.A., and G. Schwartzman. 1963. Separation and identification of sodium salts of acidic, propionic, butyric, and valeric volatile fatty acids by paper chromatography. J. of the Assoc. of Official Agricultural Chemists 46(3): 483-485.
- (556) Ziegler, H.F. Jr., and R.D. Seeley. 1961. The synergistic effect of calcium propionate and sorbic acid in inhibiting mold growth on partially baked rolls. Baker's Digest 35(3): 80.

## THIODIPROPIONIC ACID

### Bibliography

- (557) Bunnell, R.H., et al. 1955. Studies on encephalomalacia in the chick. III. The influence of feeding or injection of various tocopherols and other antioxidants on the incidence of encephalomalacia. Poultry Sci., 34: 1068-1075.
- (558) Epstein, S.S., et al. 1967. Toxicity of antioxidants to Tetrahymena pyriformis. J. Protozool. 14(2): 238-244.
- (559) French Patent 1 386 675. 1965. Addition of 5'-nucleotides to meat products.
- (560) Gribbins, M.F. 1945. Stabilization of fats and oils. U.S. Patent 2 457 227.
- (561) Gribbins, M.F., and H.R. Dittmar. 1951. Food antioxidants. U.S. Patent 2 564 106.
- (562) Hammond, E.G. 1970. Stabilizing milk fat with antioxidants. American Dairy Review 32(6): 40-41, 43, & 76-77.
- (563) Hill, I.M., et al. 1969. Protection of milk fat with antioxidants. Journal of Dairy Science 52(6): 888.
- (564) Larsson, E. 1943.  $\beta$ ,  $\beta'$ -Sulfinyldipropionic acid. Svensk Kem. Tid. 55: 29-35.
- (565) Leffler, M.T. 1951. Stable vitamin B<sub>12</sub> composition. U.S. Patent 2 579 679.
- (566) Lehman, A.J. 1950. Some toxicological reasons why certain chemicals may or may not be permitted as food additives. Assoc. Food & Drug Officials U.S., Quart. Bull. 14: 82-98.
- (566a) Lehman, A.J., et al. 1951. The pharmacological evaluation of antioxidants. Advan. Food Res. 3: 197-208.
- (567) McCaulley, D.F., et al. 1967. The multidetermination of antioxidants in lard. J. Assoc. Office. Anal. Chim. 50(2): 243-250.
- (568) Mori, T., et al. 1952-1953. Antioxidant for vitamin A-enriched margarine. II. J. Japan. Soc. Food Nutrition (Eiyo to Shokuryo) 5: 180-182.
- (569) O'Leary, D.K. 1950. Stabilized milk products. U.S. Patent 2 497 320.
- (570) O'Leary, D.K. 1947. Stabilisiertes Milchprodukt. Canada Patent 487 592
- (571) Sahasrabudhe, M.B., et al. 1964. Inhibition of tumor growth with antimetabolites of hexose monophosphate pathway intermediates. Acta Unio Intern. Contra Cancrum 20(1-2): 221-225.

\* \*

- (572) Schwab, A. W., et al. 1953. The flavor problem of soybean oil. XIII. Sulfur coordination compounds effective in edible-oil stabilization. J. Am. Oil Chemists' Soc. 30: 413-417.
- (573) Stoloff, L. S., et al. 1948. Curb mackerel fillet rancidity with new dip-and-coat technique. Food Inds. 20: 1130-1132.
- (574) Tamsma, A., et al. 1963. Factors related to the flavor stability during storage of foam-dried whole milk. III. Effect of antioxidants. Jour. Dairy Sci. 46(2): 114-119.
- \* (575) Telford, I. R., et al. 1962. Fetal resorption in the rat as influenced by certain antioxidants. Amer. Jour. Anat. 110(1): 29-36.
- (576) Thompson, J. W., and E. R. Sherwin. 1966. Investigation of antioxidants for polyunsaturated edible oils. J. Amer. Oil Chemists' Soc. 43(12): 683-686.
- (577) van der Heide, R. F., et al. 1965. Identification of antioxidants in plastics. Chem. Weekblad 61(38): 440-443.
- (578) von Riesen, V. L. 1963. Schwefelwasserstoffbildung durch *Pseudomonas aeruginosa*. 2. Mitt. Qualitative Substratuntersuchung. J. Bacteriol. 85: 248-249.
- (579) West German Patent Application 1 813 015. 1969. Antioxidant for animal and vegetable fats and oils.
- (580) Yoshikawa, K., and M. Nonaka. 1953. Antioxidants for lard and soap. Yushi Kogaku Kyôkaishi (J. Oil Chemists' Soc., Japan) 2: 186-190.

## DILAURYL THIODIPROPIONIC ACID

### Bibliography

- (581) Berndt, G., and W. Schilling. 1966. Fettstabilisation mit Antioxydantien und einem Extrakt aus *Aspergillus oryzae*. *Seifen-Ole-Fette-Wachse* 92: 81-83.
- (582) Epstein, S.S., et al. 1967. Toxicity of antioxidants to *Tetrahymena pyriformis*. *J. Protozool.* 14(2): 238-244.
- \* (583) Fazio, T., et al. 1968. Oils, fats, and waxes. Determination of DLTDP and other antioxidants by a modified sublimation technique. *J. Assoc. Offic. Anal. Chem.* 51(1): 17-20.
- (584) Lehman, A.J. 1950. Some toxicological reasons why certain chemicals may or may not be permitted as food additives. *Assoc. Food & Drug Officials U.S., Quart. Bull.* 14: 82-98.
- (585) Tamsma, A., et al. 1963. Factors relating to the flavor stability during storage of foam-dried whole milk. III Effect of antioxidants. *J. of Dairy Science* 46(2): 114-119.
- (586) Tarr, H. L. A. 1947. Control of rancidity in fish flesh. I. Chemical antioxidants. *Jour. Fish. Res. Bd. Canada* 7(3): 137-154.
- (587) Thompson, J.W., and E.R. Sherwin. 1966. Investigation of antioxidants for polyunsaturated edible oils. *J. Amer. Oil Chemists' Soc.* 43(12): 683-686.
- (588) van der Heide, R. F., et al. 1965. Identification of antioxidants in plastics. *Chem. Weekblad* 61(38): 440-443.